

MEASURING THE RANGE OF AN ADDITIVE LÉVY PROCESS

BY DAVAR KHOSHNEVISAN,¹ YIMIN XIAO¹ AND YUQUAN ZHONG²

University of Utah, Michigan State University and Academia Sinica

The primary goal of this paper is to study the range of the random field $X(t) = \sum_{j=1}^N X_j(t_j)$, where X_1, \dots, X_N are independent Lévy processes in \mathbb{R}^d .

To cite a typical result of this paper, let us suppose that Ψ_i denotes the Lévy exponent of X_i for each $i = 1, \dots, N$. Then, under certain mild conditions, we show that a necessary and sufficient condition for $X(\mathbb{R}_+^N)$ to have positive d -dimensional Lebesgue measure is the integrability of the function $\mathbb{R}^d \ni \xi \mapsto \prod_{j=1}^N \operatorname{Re}\{1 + \Psi_j(\xi)\}^{-1}$. This extends a celebrated result of Kesten and of Bretagnolle in the one-parameter setting. Furthermore, we show that the existence of square integrable local times is yet another equivalent condition for the mentioned integrability criterion. This extends a theorem of Hawkes to the present random fields setting and completes the analysis of local times for additive Lévy processes initiated in a companion by paper Khoshnevisan, Xiao and Zhong.

Contents

1. Introduction
 2. Images and local times
 3. Some calculations
 4. Proof of Theorem 2.1: upper bound
 5. Proofs of Theorems 1.1, 1.3 and 1.5
 6. Convolution-based energies
 7. Bessel–Riesz capacities
 8. Proof of Theorem 1.6
 9. Concluding remarks
- Appendix

1. Introduction. An N -parameter d -dimensional random field $X = \{X(t); t \in \mathbb{R}_+^N\}$ is an *additive Lévy process* if X has the following pathwise decomposition:

$$X(t) = X_1(t_1) + \dots + X_N(t_N) \quad \forall t \in \mathbb{R}_+^N,$$

where X_1, \dots, X_N are independent classical Lévy processes on \mathbb{R}^d . Using tensor notation, we will often write $X = X_1 \oplus \dots \oplus X_N$ for brevity, and we will always

Received December 2001; revised May 2002.

¹Supported in part by NSF and NATO grants.

²Supported by the National Natural Science Foundation of China.

AMS 2000 subject classifications. 60G60, 60J55, 60J45.

Key words and phrases. Additive Lévy processes, strictly stable processes, capacity, energy, local times, Hausdorff dimension.

assume that $X_j(0) = 0$ for all $j = 1, \dots, N$. Finally, if Ψ_1, \dots, Ψ_N denote the Lévy exponents of X_1, \dots, X_N , respectively, we define the Lévy exponent of X to be $\Psi = (\Psi_1, \dots, \Psi_N)$. See the companion paper [Khoshnevisan, Xiao and Zhong (2002)] for more detailed historical information, as well as a number of collected facts about additive Lévy processes.

The following question is the starting point of our investigation:

$$(1.1) \quad \text{“When can the range of } X \text{ have positive Lebesgue measure?”}$$

In the one-parameter setting, that is, when $N = 1$, this question has a long history as well as the following remarkable answer, discovered by Bretagnolle (1971) and Kesten (1969): If λ_d denotes Lebesgue measure in \mathbb{R}^d ,

$$(1.2) \quad \mathbb{E}\{\lambda_d(X(\mathbb{R}_+))\} > 0 \iff \int_{\mathbb{R}^d} \operatorname{Re}\left(\frac{1}{1 + \Psi(\xi)}\right) d\xi < +\infty,$$

where Ψ denotes the Lévy exponent of X , and $\operatorname{Re} z$ denotes the real part of $z \in \mathbb{C}$. In the sequel, the imaginary part and the conjugate of z will be denoted by $\operatorname{Im} z$ and \bar{z} , respectively.

The primary objective of this paper is to answer question (1.1) for the range of an additive Lévy process $X = \{X(t); t \in \mathbb{R}_+^N\}$. It is quite standard to show that $\prod_{j=1}^N \operatorname{Re}\{1 + \Psi_j\}^{-1} \in L^1(\mathbb{R}^d)$ is a sufficient condition for $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} > 0$. The converse is much more difficult to prove, and we have succeeded in doing so as long as there exists a positive constant $\vartheta > 0$ such that

$$(1.3) \quad \operatorname{Re}\left(\prod_{j=1}^N \frac{1}{1 + \Psi_j(\xi)}\right) \geq \vartheta \prod_{j=1}^N \operatorname{Re}\left(\frac{1}{1 + \Psi_j(\xi)}\right);$$

see Theorem 1.1 below. We note that, when $N = 1$, condition (1.3) holds vacuously with $\vartheta = 1$.

Among other things, we will show in this paper that, when (1.3) holds, the proper setting for the analysis of question (1.1) is potential theory and its various connections to the random field X , as well as energy that we will describe below. Various aspects of the potential theory of multiparameter processes have been treated in Evans (1987a, b), Fitzsimmons and Salisbury (1989), Hirsch (1995), Hirsch and Song (1995a, b) and Khoshnevisan and Xiao (2002a).

As we mentioned earlier, we propose to derive the following multiparameter version of (1.2). It will be a consequence of some of the later results of this article.

THEOREM 1.1. *Let X be an additive Lévy process in \mathbb{R}^d with Lévy exponent (Ψ_1, \dots, Ψ_N) , and suppose that condition (1.3) holds. Then*

$$\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} > 0 \iff \int_{\mathbb{R}^d} \prod_{j=1}^N \operatorname{Re}\left(\frac{1}{1 + \Psi_j(\xi)}\right) d\xi < +\infty.$$

REMARK 1.2. We record the fact that condition (1.3) is only needed for proving the direction “ \Rightarrow .” We also mention the fact that, under the conditions of Proposition 6.5 below, $\lambda_d(X(\mathbb{R}_+^N))$ is almost surely equal to $+\infty$ (resp. 0), if $\int_{\mathbb{R}^d} \prod_{j=1}^N \operatorname{Re}\{1 + \Psi_j(\xi)\}^{-1} d\xi$ is finite (resp. infinite).

Theorem 1.1 has the following equivalent formulation which addresses existence questions for the local times of the companion paper [Khoshnevisan, Xiao and Zhong (2002)]. When $N = 1$, it is a well-known theorem of Hawkes (1986).

THEOREM 1.3. *Let X be an additive Lévy process in \mathbb{R}^d that satisfies condition (1.3). Then X has square integrable local times if and only if $\prod_{j=1}^N \operatorname{Re}\{1 + \Psi_j\}^{-1}$ is integrable in \mathbb{R}^d , where (Ψ_1, \dots, Ψ_N) is the Lévy exponent of X .*

We have already mentioned that Theorem 1.3 is an equivalent probabilistic interpretation of Theorem 1.1. However, in fact, our formulation of Theorem 1.3 lies at the heart of our proof of Theorem 1.1 and its further refinements (cf. Theorem 2.1).

REMARK 1.4. When $N = 1$, condition (1.3) always holds with $\vartheta = 1$. Hence, our theorems extend those of Bretagnolle (1971), Kesten (1969) and Hawkes (1986).

In general, any additive Lévy process X with Lévy exponent $\Psi = (\Psi_1, \dots, \Psi_N)$ induces an energy form \mathcal{E}_Ψ that can be described as follows: For all finite measures μ on \mathbb{R}^d , and/or all integrable functions $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$(1.4) \quad \mathcal{E}_\Psi(\mu) = (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^2 \prod_{j=1}^N \operatorname{Re}\left(\frac{1}{1 + \Psi_j(\xi)}\right) d\xi,$$

where $\widehat{\cdot}$ denotes the Fourier transform normalized as $\widehat{f}(\xi) = \int_{\mathbb{R}^d} e^{i\xi \cdot x} f(x) dx$ [$f \in L^1(\mathbb{R}^d)$].

Frequently, we may refer to “the energy” of a measure (or function) in the context of an additive Lévy process X without explicitly mentioning its dependence on the Lévy exponent of X . This makes for a simpler presentation and should not cause ambiguities.

Having introduced energies, we can present a key result of this paper. When $N = 1$, it can be found in Bertoin [(1996), page 60].

THEOREM 1.5. *Consider any d -dimensional additive Lévy process X whose Lévy exponent Ψ satisfies (1.3). Then, given any nonrandom compact set $F \subset \mathbb{R}^d$, $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N) \oplus F)\} > 0$ if and only if F carries a finite measure of finite energy.*

We have adopted the notation that, for all sets A and B , $A \oplus B = \{a + b; a \in A, b \in B\}$. This should not be confused with our tensor notation for $X = X_1 \oplus \dots \oplus X_N$.

Note, in particular, that if we choose F to be a singleton in Theorem 1.5, we immediately obtain Theorem 1.1.

Next, we apply Theorem 1.5 to compute the Hausdorff dimension of the range of an arbitrary additive Lévy process.

THEOREM 1.6. *Given an additive Lévy process X in \mathbb{R}^d with Lévy exponent (Ψ_1, \dots, Ψ_N) that satisfies (1.3),*

$$\dim(X(\mathbb{R}_+^N)) = d - \eta, \quad \mathbb{P}\text{-a.s.},$$

where

$$\eta = \sup \left\{ \alpha > 0 : \int_{\xi \in \mathbb{R}^d : \|\xi\| > 1} \prod_{j=1}^N \operatorname{Re} \left(\frac{1}{1 + \Psi_j(\xi)} \right) \frac{d\xi}{\|\xi\|^\alpha} = +\infty \right\}.$$

Here, $\dim(\cdot)$ denotes Hausdorff dimension, $\sup \emptyset = 0$, and $\|\cdot\|$ denotes the Euclidean ℓ^2 -norm.

REMARK 1.7. It can be checked directly that

$$\eta = \inf \left\{ \alpha > 0 : \int_{\xi \in \mathbb{R}^d : \|\xi\| > 1} \prod_{j=1}^N \operatorname{Re} \left(\frac{1}{1 + \Psi_j(\xi)} \right) \frac{d\xi}{\|\xi\|^\alpha} < +\infty \right\}.$$

Furthermore, one always has $\eta \leq d$.

When $N = 1$, that is, when X is an ordinary Lévy process in \mathbb{R}^d , Pruitt (1969) has shown that the Hausdorff dimension of the range of X is

$$\gamma = \sup \left\{ \alpha \geq 0 : \limsup_{r \rightarrow 0} r^{-\alpha} \int_0^1 \mathbb{P}\{|X(t)| \leq r\} dt < +\infty \right\}.$$

In general, this computation is not satisfying, since the above lim sup is not easy to evaluate. Pruitt [(1969), Theorem 5] addresses this issue by verifying the following estimate for γ in terms of the Lévy exponent Ψ of X :

$$\gamma \geq \sup \left\{ \alpha < d : \int_{\|\xi\| \geq 1} \frac{1}{|\Psi(\xi)|} \frac{d\xi}{\|\xi\|^{d-\alpha}} < +\infty \right\}.$$

Moreover, it is shown there that if, in addition, $\operatorname{Re} \Psi(\xi) \geq 2 \log \|\xi\|$ (for all $\|\xi\|$ large), then

$$\gamma = \sup \left\{ \alpha < d : \int_{\mathbb{R}^d} \operatorname{Re} \left(\frac{1 - e^{-\Psi(\xi)}}{\Psi(\xi)} \right) \frac{d\xi}{\|\xi\|^{d-\alpha}} < +\infty \right\}.$$

See Fristedt [(1974), pages 377–378] for further discussions on Pruitt’s work in this area. Our Theorem 1.6 readily implies the following representation for the index γ that holds under no restrictions. To the best of our knowledge, it is new.

COROLLARY 1.8. *If X denotes a Lévy process in \mathbb{R}^d with Lévy exponent Ψ ,*

$$\gamma = \sup \left\{ \alpha < d : \int_{\xi \in \mathbb{R}^d : \|\xi\| > 1} \operatorname{Re} \left(\frac{1}{1 + \Psi(\xi)} \right) \frac{d\xi}{\|\xi\|^{d-\alpha}} < +\infty \right\}.$$

REMARK 1.9. To paraphrase Kesten [(1969), page 7], (1.2) has the somewhat unexpected consequence that the range of a Lévy process $\{X(t); t \geq 0\}$ has a better chance of having positive Lebesgue’s measure than the range of its symmetrization $\{X(t) - X'(t); t \geq 0\}$, where X' is an independent copy of X . Thanks to Corollary 1.8, this qualitative statement has a *quantitative* version. Namely, with probability 1, $\dim(X(\mathbb{R}_+)) \geq \dim(Y(\mathbb{R}_+))$, where $Y(t) = X(t) - X'(t)$ is the symmetrization of X . To prove this, one need only note that $\operatorname{Re}\{1 + \Psi\}^{-1} \leq \{1 + \operatorname{Re} \Psi\}^{-1}$, pointwise.

REMARK 1.10. The preceding remark can be adapted to show that for any additive Lévy process $\{X(t); t \in \mathbb{R}_+^N\}$ that satisfies (1.3),

$$(1.5) \quad \dim(X(\mathbb{R}_+^N)) \geq \dim(Y(\mathbb{R}_+^N)) \quad \text{a.s.}$$

Here, Y is the symmetrization of X defined by $Y(t) = X(t) - X'(t)$, where X' is an independent copy of X . To verify the displayed inequality, we first note that Y also satisfies (1.3) (cf. Example 1.16). Thus, our claim follows from Theorem 1.6 and the elementary pointwise inequality: $\prod_{j=1}^N \operatorname{Re}\{1 + \Psi_j\}^{-1} \leq \prod_{j=1}^N \{1 + \operatorname{Re} \Psi_j\}^{-1}$. Furthermore, we note that the strict inequality in (1.5) may hold even for $N = 1$; see Pruitt [(1969), Section 4] for an example. This example was also noticed by Taylor [(1973), page 401], but there was a minor error in his statement on line –3: “smaller” should be “larger.” It is worthwhile to point out that Hawkes (1974), by modifying the construction of Pruitt (1969), defined another Lévy process X in \mathbb{R} such that its range has positive one-dimensional Lebesgue measure, while the Hausdorff dimension of the range of its symmetrization is strictly smaller than 1.

REMARK 1.11. There are several interesting “indices” for Lévy processes, one of which is the index γ mentioned earlier. These indices arise when studying various properties of the sample paths of Lévy processes and include the *upper index* β , the *lower indices* β' and β'' [Blumenthal and Gettoor (1961)] and the index γ' [Hendricks (1983)]. Rather than reintroducing these indices, we only mention that

$$0 \leq \beta' \wedge d \leq \gamma \leq \gamma' \leq \beta \wedge d.$$

Pruitt and Taylor (1996) discuss some open problems regarding these indices.

The following is an outline of the paper: In Section 2 (Theorem 2.1) we state a complete characterization of all compact sets $E \subset \mathbb{R}_+^N$ for which the stochastic

image $X(E)$ can have positive Lebesgue measure. After establishing a number of preparatory lemmas about the semigroup and the resolvent of an additive Lévy process in Section 3, we complete our proof of Theorem 2.1 in Section 4. Our proofs of Theorems 1.1, 1.3 and 1.5 can be found in Section 5. In Section 6 we briefly discuss some of the existing connections between the energy $\mathcal{E}_\Psi(\mu)$ —introduced in (1.4)—and classical convolution-based energy forms. In Section 7 we utilize additive Lévy processes to describe a probabilistic interpretation of all sets of positive α -dimensional Bessel–Riesz capacity where $\alpha \geq 0$ is arbitrary. This probabilistic representation is used in Section 8, where Theorem 1.6 is derived. In Section 9, we have stated some remaining open problems.

Since condition (1.3) will play an important role in our arguments, we end this section with some examples of additive Lévy processes that satisfy (1.3).

EXAMPLE 1.12. Consider the following condition:

(1.6) At least $N - 1$ of the Lévy processes X_1, \dots, X_N are symmetric.

By using induction, we can see that the above implies that

$$(1.7) \quad \operatorname{Re} \left(\prod_{j=1}^N \frac{1}{1 + \Psi_j(\xi)} \right) = \prod_{j=1}^N \operatorname{Re} \left(\frac{1}{1 + \Psi_j(\xi)} \right).$$

In particular, condition (1.6) implies (1.3) with $\vartheta = 1$. [It may help to recall that an ordinary Lévy process Y is *symmetric* if $Y(1)$ and $-Y(1)$ have the same distribution.]

EXAMPLE 1.13. Consider a two-parameter additive Lévy process $\mathbb{R}_+^2 \ni t \mapsto X_1(t_1) + X_2(t_2)$, where X_1 and X_2 are i.i.d. Lévy processes on \mathbb{R}^d with exponent Ψ_1 each. Then it is possible to directly check that condition (1.3) holds if and only if

$$(1.8) \quad \exists \delta \in (0, 1) : \forall \xi \in \mathbb{R}^d, \quad |\operatorname{Im} \Psi_1(\xi)| \leq \delta(1 + \operatorname{Re} \Psi_1(\xi)).$$

This is a kind of sector condition on Ψ_1 .

EXAMPLE 1.14. Suppose X_1 and X_2 are independent Lévy processes on \mathbb{R}^d , and with Lévy exponents Ψ_1 and Ψ_2 , respectively. Then one can check directly that the two-parameter additive Lévy process $\mathbb{R}_+^2 \ni t \mapsto X_1(t_1) - X_2(t_2)$ satisfies condition (1.3) as long as

$$\operatorname{Im} \Psi_1(\xi) \cdot \operatorname{Im} \Psi_2(\xi) \geq 0 \quad \forall \xi \in \mathbb{R}^d.$$

In particular, if X_1 and X_2 are i.i.d., condition (1.3) *always* holds for the process $t \mapsto X_1(t_1) - X_2(t_2)$. This process arises in studying the self-intersections of Lévy processes.

EXAMPLE 1.15. Suppose $N > 2$, and consider the N -parameter additive Lévy process $X = X_1 \oplus \cdots \oplus X_N$, where X_1, \dots, X_N are i.i.d. Lévy processes on \mathbb{R}^d , all with the same Lévy exponent Ψ_1 . Writing in polar coordinates, we have $\{1 + \Psi_1(\xi)\}^{-1} = r e^{i\theta}$, where $r = |1 + \Psi_1(\xi)|^{-1}$, and $\cos(\theta) = |1 + \Psi_1(\xi)|^{-1} \{1 + \operatorname{Re} \Psi_1(\xi)\}$. According to Taylor’s formula,

$$\cos(N\theta) \geq 1 - \frac{1}{2}N^2\theta^2 \geq \frac{1}{2} \geq \frac{1}{2}(\cos \theta)^N,$$

as long as $|\theta| \leq N^{-1}$. Consequently, $|\theta| \leq N^{-1}$ implies condition (1.3). Equivalently,

$$|\operatorname{Im} \Psi_1(\xi)| \leq \tan\left(\frac{1}{N}\right) |1 + \operatorname{Re} \Psi_1(\xi)| \quad \forall \xi \in \mathbb{R}^d$$

implies condition (1.3).

EXAMPLE 1.16 (Symmetrization). Suppose $X_1, \dots, X_k; \bar{X}_1, \dots, \bar{X}_k$ are independent Lévy processes on \mathbb{R}^d with Lévy exponents $\Psi_1, \dots, \Psi_k; \bar{\Psi}_1, \dots, \bar{\Psi}_k$, respectively, where $\bar{\Psi}_\ell$ denotes the complex conjugate of Ψ_ℓ ($\ell = 1, \dots, k$). Then the additive Lévy process V satisfies condition (1.3) with $\vartheta = 1$, where

$$V = X_1 \oplus \cdots \oplus X_k \oplus \bar{X}_1 \oplus \cdots \oplus \bar{X}_k.$$

If, in addition, Y is an arbitrary \mathbb{R}^d -valued Lévy process that is independent of V , the additive Lévy process $V \oplus Y$ also satisfies (1.3) with $\vartheta = 1$.

2. Images and local times. Throughout, we let $X = X_1 \oplus \cdots \oplus X_N$ denote a d -dimensional additive Lévy process with Lévy exponent $\Psi = (\Psi_1, \dots, \Psi_N)$. In this section we seek to find a general condition that guarantees that the image $X(E)$ of a compact set $E \subset \mathbb{R}_+^N$ can have positive Lebesgue measure. Under regularity conditions on X , this was achieved in Khoshnevisan and Xiao (2002a, b). Our goal here is to find conditions for the positivity of the image $\lambda_d(X(E))$ that hold quite generally.

Any finite measure μ on \mathbb{R}_+^N defines an *occupation measure* \mathbb{O}_μ on \mathbb{R}^d via the prescription

$$(2.1) \quad \mathbb{O}_\mu(A) = \int_{\mathbb{R}_+^N} \mathbb{1}_A(X(s)) \mu(ds), \quad A \in \mathcal{B}(\mathbb{R}^d),$$

where $\mathbb{1}_A(\cdot)$ is the indicator function of A and $\mathcal{B}(\mathbb{R}^d)$ is the Borel σ -field of \mathbb{R}^d .

We take the distribution approach to measures. In particular, we tacitly identify the preceding random measure with the random linear operator \mathbb{O}_μ defined as

$$\mathbb{O}_\mu(f) = \int_{\mathbb{R}_+^N} f(X(s)) \mu(ds).$$

The following result is the main inequality of this section, where $\mathcal{P}(E)$ is the collection of all probability measures on E .

THEOREM 2.1. *For all compact sets $E \subset \mathbb{R}_+^N$,*

$$\begin{aligned} & \left[(2\pi)^{-d} \inf_{\mu \in \mathcal{P}(E)} \mathbb{E} \left\{ \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2 \right\} \right]^{-1} \\ & \leq \mathbb{E} \{ \lambda_d(X(E)) \} \\ & \leq 16^N \left[(2\pi)^{-d} \inf_{\mu \in \mathcal{P}(E)} \mathbb{E} \left\{ \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2 \right\} \right]^{-1}. \end{aligned}$$

REMARK 2.2. Condition (1.3) is *not* needed here.

Consequently, for $\lambda_d(X(E))$ to have positive expectation, it is necessary, as well as sufficient, that for some probability measure μ on E , the $L^2(\mathbb{R}^d)$ -norm of the Fourier transform of \mathbb{O}_μ be square integrable with respect to \mathbb{P} .

Suppose, then, that $\mathbb{E} \{ \lambda_d(X(E)) \} > 0$. Thanks to the foregoing discussion, there exists $\mu \in \mathcal{P}(E)$ such that $\|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}$ is in $L^2(\mathbb{P})$; in particular, it is finite, a.s. By Plancherel’s theorem, \mathbb{O}_μ is absolutely continuous with respect to λ_d , a.s. Let L_μ^x denote this density. In other words, $L_\mu = \{L_\mu^x; x \in \mathbb{R}^d\}$ is the process defined by the following: For all measurable functions $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$,

$$(2.2) \quad \mathbb{O}_\mu(f) = \int_{\mathbb{R}^d} f(x) L_\mu^x dx$$

[cf. (2.1)]. We can always choose a measurable version of $\mathbb{R}^d \times \Omega \ni (x, \omega) \mapsto L_\mu^x(\omega)$, which we take for granted. Of course, Ω denotes the underlying probability space. Furthermore, we can apply Plancherel’s formula, once again, to deduce that

$$(2.3) \quad \|L_\mu\|_{L^2(\mathbb{R}^d)}^2 = (2\pi)^{-d} \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2, \quad \mathbb{P}\text{-a.s.}$$

The process L_μ is the *local times of X* , under the measure μ . The above, together with Theorem 2.1, shows the following.

COROLLARY 2.3. *For $\mathbb{E} \{ \lambda_d(X(E)) \}$ to be positive, it is necessary and sufficient that there exists a probability measure μ on E , under which there are local times L_μ such that $\mathbb{E} \{ \|L_\mu\|_{L^2(\mathbb{R}^d)}^2 \}$ is finite.*

We have developed the requisite material for the lower bound (i.e., the “easy half”) in Theorem 2.1.

PROOF THEOREM 2.1 (Lower bound). Without loss of generality, we may assume that there is a probability measure μ on the Borel set E such that $\mathbb{E} \{ \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2 \} < +\infty$. Let L_μ denote the corresponding local times. It follows

from (2.2) with $f(x) = \mathbb{1}_{X(E)}(x)$ that, \mathbb{P} -a.s.,

$$\begin{aligned} 1 &= |\mathbb{O}_\mu(\mathbb{R}^d)|^2 \\ &= \left| \int_{X(E)} L_\mu^x dx \right|^2 \\ &\leq (2\pi)^{-d} \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2 \cdot \lambda_d(X(E)), \end{aligned}$$

thanks to (2.3) and to the Cauchy–Schwarz inequality. Next, recall that for all positive random variables Z , $\mathbb{E}\{Z^{-1}\} \cdot \mathbb{E}\{Z\} \geq 1$. This also follows from the Cauchy–Schwarz inequality or from Jensen’s inequality. Thus, we obtain

$$\begin{aligned} \mathbb{E}\{\lambda_d(X(E))\} &\geq (2\pi)^d \mathbb{E}\left\{ \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^{-2} \right\} \\ &\geq (2\pi)^d \left[\mathbb{E}\left\{ \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2 \right\} \right]^{-1}. \end{aligned}$$

This proves the lower bound in Theorem 2.1. \square

We conclude this section with the following analytical description of $\mathbb{E}\{\|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2\}$. Its derivation is simple, but we include it as a natural way to introduce the *associated process* \tilde{X} in (2.4) below. The remainder of Theorem 2.1 is proved in Section 4 after our presentation of Section 3, which is concerned with some calculations.

LEMMA 2.4. *For any finite measure μ on \mathbb{R}_+^N ,*

$$\begin{aligned} &\mathbb{E}\left\{ \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2 \right\} \\ &= \int_{\mathbb{R}^d} \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \exp\left(- \sum_{j=1}^N |s_j - t_j| \Psi_j(\text{sgn}(s_j - t_j)\xi) \right) \mu(ds) \mu(dt) d\xi. \end{aligned}$$

PROOF. This is an exercise in Fubini’s theorem. Indeed, by (2.1),

$$\mathbb{E}\left\{ \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2 \right\} = \int_{\mathbb{R}^d} \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \mathbb{E}\{e^{i\xi \cdot [X(s) - X(t)]}\} \mu(ds) \mu(dt) d\xi.$$

Define the N -parameter process $\tilde{X} = \{\tilde{X}(t); t \in \mathbb{R}^N\}$ by

$$(2.4) \quad \tilde{X}(t) = \sum_{j=1}^N \text{sgn}(t_j) X_j(|t_j|) \quad \forall t \in \mathbb{R}^N.$$

We emphasize that \tilde{X} is a process indexed by all of \mathbb{R}^N , and that

$$(2.5) \quad \forall s, t \in \mathbb{R}_+^N : X(t) - X(s) \text{ has the same distribution as } \tilde{X}(t - s).$$

Moreover,

$$(2.6) \quad \mathbb{E}\{e^{i\xi \cdot \tilde{X}(t)}\} = \exp\left(-\sum_{j=1}^N |t_j| \Psi_j(\text{sgn}(t_j)\xi)\right) \quad \forall \xi \in \mathbb{R}^d, t \in \mathbb{R}^N.$$

Our lemma follows. \square

3. Some calculations. Recall the associated process \tilde{X} from (2.4), and let $P = \{P_t; t \in \mathbb{R}^N\}$ be the family of operators on $L^\infty(\mathbb{R}^d)$ defined by

$$(3.1) \quad P_t f(x) = \mathbb{E}\{f(\tilde{X}(t) + x)\} \quad \forall t \in \mathbb{R}^N, f \in L^\infty(\mathbb{R}^d) \text{ and } x \in \mathbb{R}^d.$$

This is not an N -parameter semigroup; that is, it is not true that $P_{t+s} = P_t P_s$ for all $s, t \in \mathbb{R}^N$. However, each of its 2^N restrictions $\{P_t; t \in (\pm\mathbb{R})^N\}$ is an N -parameter semigroup. Let U denote the 1-potential of the family P . That is, for all $f \in L^\infty(\mathbb{R}^d)$,

$$(3.2) \quad Uf(x) = \int_{\mathbb{R}^N} \exp\left(-\sum_{j=1}^N |s_j|\right) P_s f(x) ds \quad \forall x \in \mathbb{R}^d.$$

We will also need the following potential operator:

$$(3.3) \quad U_+ f(x) = \int_{\mathbb{R}_+^N} \exp\left(-\sum_{j=1}^N s_j\right) P_s f(x) ds \quad \forall x \in \mathbb{R}^d.$$

Our next lemma computes $P_t f(x)$, $Uf(x)$ and $U_+ f(x)$ in terms of the Lévy exponent of X . In light of Theorem 1.1, it shows that \tilde{X} and its 1-potential U are the “right” objects to consider. Indeed, the integral of Theorem 1.1 is nothing but $U\delta_0(0)$, where δ_a is point mass at $a \in \mathbb{R}^d$, while condition (1.3) allows us to compare $U\delta_0(0)$ and $U_+\delta_0(0)$.

LEMMA 3.1. *The operators P_t , U and U_+ are convolution operators. Moreover, if $f, \hat{f} \in L^1(\mathbb{R}^d)$, then for all $t \in \mathbb{R}^N$ and all $x \in \mathbb{R}^d$,*

$$P_t f(x) = (2\pi)^{-d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \hat{f}(-\xi) \exp\left(-\sum_{j=1}^N |t_j| \Psi_j(\text{sgn}(t_j)\xi)\right) d\xi,$$

$$Uf(x) = 2^N (2\pi)^{-d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \hat{f}(-\xi) \prod_{j=1}^N \text{Re}\left(\frac{1}{1 + \Psi_j(\xi)}\right) d\xi,$$

$$U_+ f(x) = (2\pi)^{-d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \hat{f}(-\xi) \prod_{j=1}^N \frac{1}{1 + \Psi_j(\xi)} d\xi.$$

PROOF. Temporarily let μ_t denote the distribution of $-\tilde{X}(t)$ to see that $P_t f(x) = \mu_t \star f(x)$ and $Uf(x) = \int_{\mathbb{R}^N} \exp(-\sum_{j=1}^N |t_j|) \mu_t \star f(x) dt$, where \star denotes convolution. Since $f \in L^1(\mathbb{R}^d)$, it follows from Fubini's theorem and (2.6) that

$$\begin{aligned} \widehat{P_t f}(\xi) &= \widehat{f}(\xi) \widehat{\mu_t}(\xi) \\ (3.4) \qquad &= \widehat{f}(\xi) \exp\left(-\sum_{j=1}^N |t_j| \Psi_j(-\operatorname{sgn}(t_j)\xi)\right). \end{aligned}$$

Hence, the asserted formula for $P_t f(x)$ follows from the inversion theorem of Fourier transforms, after making a change of variables. To obtain the second equation, we integrate $P_t f$, namely,

$$Uf(x) = (2\pi)^{-d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \widehat{f}(-\xi) \int_{\mathbb{R}^N} \exp\left(-\sum_{j=1}^N |s_j| [1 + \Psi_j(\operatorname{sgn}(s_j)\xi)]\right) ds d\xi.$$

On the other hand,

$$\begin{aligned} &\int_{\mathbb{R}^N} \exp\left(-\sum_{j=1}^N |s_j| [1 + \Psi_j(\operatorname{sgn}(s_j)\xi)]\right) ds \\ &= \prod_{j=1}^N \left\{ \int_0^\infty e^{-s[1+\Psi_j(\xi)]} ds + \int_0^\infty e^{-s[1+\overline{\Psi_j(\xi)}]} ds \right\} \\ &= 2^N \prod_{j=1}^N \operatorname{Re}\{1 + \Psi_j(\xi)\}^{-1}. \end{aligned}$$

The mentioned computation of $Uf(x)$ follows readily from this. Our computation of $U_+ f$ is made in like fashion, and we omit the details. \square

Throughout, we assume that the underlying sample space Ω is the collection of all paths $\omega: \mathbb{R}_+^N \rightarrow \mathbb{R}^d$ that have the form $\omega(t) = \sum_{j=1}^N \omega_j(t_j)$ for $t \in \mathbb{R}_+^N$, where ω_j is in $D_{\mathbb{R}^d}[0, \infty)$ —the usual space of \mathbb{R}^d -valued cadlag functions—for every $j = 1, \dots, N$. The space Ω inherits its Borel field from the Skorohod topology on $D_{\mathbb{R}^d}[0, \infty)$ in a standard way. The additive Lévy process X of the Introduction is in *canonical form* if

$$X(t)(\omega) = \omega(t) \qquad \forall \omega \in \Omega, t \in \mathbb{R}_+^N.$$

Since we are only interested in distributional results about X , we can assume, with no loss in generality, that it is in canonical form under a fixed probability measure, denoted by \mathbb{P} . This is a standard result and we will not dwell on it here. Henceforth, the canonical form of X is tacitly assumed. We will also assume, with no further

mention, that $\Psi = (\Psi_1, \dots, \Psi_N)$ is the Lévy exponent of the additive process X . We note for future reference that this is equivalent to

$$(3.5) \quad \mathbb{E}\{e^{i\xi \cdot X(t)}\} = e^{-t \cdot \Psi(\xi)} \quad \forall t \in \mathbb{R}_+^N, \xi \in \mathbb{R}^d.$$

In agreement with the notation of classical Lévy processes, we define \mathbb{P}_x to be the law of $x + X$ for any $x \in \mathbb{R}^d$, and let \mathbb{E}_x be the corresponding expectation operator. To be precise, we define, for all Borel sets $A \subset \Omega$,

$$\mathbb{P}_x\{\omega \in \Omega : \omega \in A\} = \mathbb{P}\{\omega \in \Omega : x + \omega \in A\},$$

where $(x + \omega)(t) = x + \omega(t)$ for all $\omega \in \Omega$ and $t \in \mathbb{R}_+^N$. This allows us also to define a sigma-finite measure \mathbb{P}_{λ_d} , and a corresponding integration (or expectation) operator, \mathbb{E}_{λ_d} , via

$$\begin{aligned} \mathbb{P}_{\lambda_d}\{A\} &= \int_{\mathbb{R}^d} \mathbb{P}_x\{A\} dx & \forall A \subset \mathbb{R}^d \text{ Borel,} \\ \mathbb{E}_{\lambda_d}\{Z\} &= \int_{\mathbb{R}^d} \mathbb{E}_x\{Z\} dx & \forall Z : \Omega \rightarrow \mathbb{R}_+, \text{ measurable.} \end{aligned}$$

The last line holds for a larger class of random variables Z by standard monotone class arguments.

Let $\Pi = \{1, \dots, N\}$, and for all $A \subseteq \Pi$ define the partial order $\preceq_{(A)}$ on \mathbb{R}^N by

$$s \overset{(A)}{\preceq} t \iff \begin{cases} s_i \leq t_i, & \text{for all } i \in A, \\ s_i \geq t_i, & \text{for all } i \in A^c. \end{cases}$$

We may also write $t \succ_{(A)} s$ for $s \preceq_{(A)} t$. We note that, used in conjunction, the partial orders $\{\preceq_{(A)}; A \subseteq \Pi\}$ totally order \mathbb{R}^N in the sense that, for all $s, t \in \mathbb{R}^N$, we can find $A \subseteq \Pi$ such that $s \preceq_{(A)} t$. We will use this simple fact several times. The final piece of notation is that of filtrations in each partial order $\preceq_{(A)}$. Namely, we define $\mathcal{F}^A(t)$ to be the sigma-field generated by $\{X(r); r \preceq_{(A)} t\}$. We can, and will, assume that each $\mathcal{F}^A(t)$ is \mathbb{P}_x -complete for all $x \in \mathbb{R}^d$, and each \mathcal{F}^A is $\preceq_{(A)}$ -right continuous. The latter means that, for all $t \in \mathbb{R}_+^N$, $\mathcal{F}^A(t) = \bigcap_{s \succ_{(A)} t} \mathcal{F}^A(s)$.

The following key fact is borrowed from Khoshnevisan and Xiao (2002a), which we reproduce for the sake of completeness.

PROPOSITION 3.2 (The Markov property). *Suppose that $A \subseteq \Pi$ and that $s \preceq_{(A)} t$, where s and t are both in \mathbb{R}_+^N . Then, for any measurable function $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$,*

$$\mathbb{E}_{\lambda_d}\{f(X(t)) | \mathcal{F}^A(s)\} = P_{t-s} f(X(s)),$$

\mathbb{P}_{λ_d} -a.s., where P_t is defined in (3.1).

REMARK 3.3. This is not generally true under \mathbb{P}_x . Also note that conditional expectations under the sigma-finite measure \mathbb{P}_{λ_d} are defined in exactly the same manner as those with respect to probability measures.

PROOF OF PROPOSITION 3.2. Consider measurable functions $f, g, h_1, \dots, h_m : \mathbb{R}^d \rightarrow \mathbb{R}_+$ and times $t, s, s^1, \dots, s^m \in \mathbb{R}_+^N$ such that $t \succ_{(A)} s \succ_{(A)} s^j$ for all $j = 1, \dots, m$. Then, since the X_j 's are independent from one another, and by appealing to the independent-increments property of each X_j , we deduce

$$\begin{aligned} & \mathbb{E}_{\lambda_d} \left\{ f(X(t)) \cdot g(X(s)) \cdot \prod_{j=1}^m h_j(X(s^j)) \right\} \\ &= \int_{\mathbb{R}^d} \mathbb{E} \left\{ f(X(t) + x) \cdot g(X(s) + x) \cdot \prod_{j=1}^m h_j(X(s^j) + x) \right\} dx \\ &= \int_{\mathbb{R}^d} \mathbb{E} \{ f(X(t) - X(s) + y) \} \mathbb{E} \left\{ \prod_{j=1}^m h_j(X(s^j) - X(s) + y) \right\} g(y) dy. \end{aligned}$$

Thanks to (2.4), (2.5) and (3.1), the first term under the integral equals $P_{t-s} f(y)$. Noting that, under the measure \mathbb{P}_{λ_d} , the distribution of $X(s)$ is λ_d , we see that the desired result follows. \square

LEMMA 3.4. Suppose $f, g : \mathbb{R}^d \rightarrow \mathbb{R}$ are in $L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ and their Fourier transforms are in $L^1(\mathbb{R}^d)$. Then, for all $s, t \in \mathbb{R}_+^N$,

$$\begin{aligned} & \mathbb{E}_{\lambda_d} \{ f(X(s)) g(X(t)) \} \\ &= (2\pi)^{-d} \int_{\mathbb{R}^d} \widehat{f}(\xi) \overline{\widehat{g}(\xi)} \exp\left(-\sum_{j=1}^N |t_j - s_j| \Psi_j(\text{sgn}(t_j - s_j)\xi)\right) d\xi. \end{aligned}$$

PROOF. Find $A \subseteq \Pi$ such that $s \preccurlyeq_{(A)} t$. Then

$$\begin{aligned} \mathbb{E}_{\lambda_d} \{ f(X(s)) g(X(t)) \} &= \mathbb{E}_{\lambda_d} \{ f(X(s)) \mathbb{E}_{\lambda_d} \{ g(X(t)) | \mathcal{F}^A(s) \} \} \\ &= \mathbb{E}_{\lambda_d} \{ f(X(s)) P_{t-s} g(X(s)) \}, \end{aligned}$$

thanks to Proposition 3.2. Since the \mathbb{P}_{λ_d} -distribution of $X(s)$ is λ_d for any $s \in \mathbb{R}_+^N$ and both $f, g \in L^2(\mathbb{R}^d)$, we can deduce the following from Plancherel's formula:

$$\begin{aligned} \mathbb{E}_{\lambda_d} \{ f(X(s)) g(X(t)) \} &= \int_{\mathbb{R}^d} f(x) P_{t-s} g(x) dx \\ &= (2\pi)^{-d} \int_{\mathbb{R}^d} \widehat{f}(\xi) \overline{\widehat{P_{t-s} g}(\xi)} d\xi. \end{aligned}$$

Our lemma follows from (3.4). \square

Next, we recall the occupation measures \mathbb{O}_μ from (2.1). The following is a function analogue of Lemma 2.4.

LEMMA 3.5. *For all $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ in $L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ such that $\widehat{f} \in L^1(\mathbb{R}^d)$, and for all probability measures μ on \mathbb{R}_+^N ,*

$$\mathbb{E}_{\lambda_d}\{|\mathbb{O}_\mu(f)|^2\} = (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{f}(\xi)|^2 Q_\mu(\xi) d\xi,$$

where

$$(3.6) \quad Q_\mu(\xi) = \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \exp\left(-\sum_{j=1}^N |t_j - s_j| \Psi_j(\text{sgn}(t_j - s_j)\xi)\right) \mu(ds) \mu(dt).$$

PROOF. This follows from Lemma 3.4 and Fubini’s theorem, once we verify that the function Q_μ is nonnegative. On the other hand,

$$(3.7) \quad Q_\mu(\xi) = \mathbb{E}\left\{\left|\int_{\mathbb{R}_+^N} e^{i\xi \cdot X(t)} \mu(dt)\right|^2\right\} \quad \forall \xi \in \mathbb{R}^d,$$

thanks to (2.5) and (2.6). This implies the pointwise positivity of Q_μ , thus concluding our proof. \square

4. Proof of Theorem 2.1: upper bound. In Section 2 we proved the easy half (i.e., the lower bound) of Theorem 2.1. We now use the results of the previous section to derive the hard half of Theorem 2.1.

For all measurable $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$, all probability measures μ on \mathbb{R}_+^N and all $A \subseteq \Pi$, define the process $\mathcal{M}_\mu^A f$ by

$$(4.1) \quad \mathcal{M}_\mu^A f(t) = \mathbb{E}_{\lambda_d}\{\mathbb{O}_\mu(f) | \mathcal{F}^A(t)\} \quad \forall t \in \mathbb{R}_+^N.$$

LEMMA 4.1. *Suppose $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ is in $L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$, and $\widehat{f} \in L^1(\mathbb{R}^d)$, $A \subseteq \Pi$, and μ is a probability measure on \mathbb{R}_+^N . Then, recalling (3.6), we have*

$$\begin{aligned} \mathbb{E}_{\lambda_d}\{\mathcal{M}_\mu^A f(t)\} &= \int_{\mathbb{R}^d} f(x) dx \quad \forall \xi \in \mathbb{R}^d, \\ \sup_{t \in \mathbb{R}_+^N} \mathbb{E}_{\lambda_d}\{|\mathcal{M}_\mu^A f(t)|^2\} &\leq (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{f}(\xi)|^2 Q_\mu(\xi) d\xi. \end{aligned}$$

PROOF. The \mathbb{P}_{λ_d} -expectation of $\mathcal{M}_\mu^A f(t)$ follows immediately from Fubini’s theorem, and the elementary fact that $\mathbb{E}_{\lambda_d}\{f(X(t))\} = \int_{\mathbb{R}^d} f(x) dx$. For the second identity, we note that, by the Cauchy–Schwarz inequality for conditional expectation under \mathbb{P}_{λ_d} , for all $t \in \mathbb{R}_+^N$, $\mathbb{E}_{\lambda_d}\{|\mathcal{M}_\mu^A f(t)|^2\} \leq \mathbb{E}_{\lambda_d}\{|\mathbb{O}_\mu(f)|^2\}$. The lemma follows from Lemma 3.5. \square

LEMMA 4.2. For all $s \in \mathbb{R}_+^N$, all measurable functions $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ in $L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ with $\widehat{f} \in L^1(\mathbb{R}^d)$, and for any probability measure μ on \mathbb{R}_+^N , \mathbb{P}_{λ_d} -almost surely, the following holds:

$$\sum_{A \subseteq \Pi} \mathcal{M}_\mu^A f(s) \geq \int_{\mathbb{R}_+^N} P_{t-s} f(X(s)) \mu(dt).$$

Moreover,

$$\mathbb{E}_{\lambda_d} \left\{ \sup_{s \in \mathbb{Q}_+^N} |\mathcal{M}_\mu^A f(s)|^2 \right\} \leq 4^N (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{f}(\xi)|^2 Q_\mu(\xi) d\xi.$$

REMARK 4.3. Since our filtrations satisfy the “usual conditions,” one can show that, when f is bounded, say, $\mathcal{M}_\mu^A f$ has a $\preceq_{(A)}$ -right continuous modification. Consequently, for this modification, the former inequality holds almost surely, where the null set in question is independent of $t \in \mathbb{R}_+^N$. See Bakry (1979) for a version of such a regularity result.

PROOF OF LEMMA 4.2. For the first expression, we note from (4.1) that since $f \geq 0$,

$$\begin{aligned} \mathcal{M}_\mu^A f(s) &\geq \mathbb{E}_{\lambda_d} \left\{ \int_{t \succ_{(A)} s} f(X(t)) \mu(dt) \middle| \mathcal{F}^A(s) \right\} \\ &= \int_{t \succ_{(A)} s} P_{t-s} f(X(s)) \mu(dt), \end{aligned}$$

\mathbb{P}_{λ_d} -a.s. for any probability measure μ on \mathbb{R}^d , thanks to Proposition 3.2. Summing this over all $A \subseteq \Pi$ and recalling that, together, the $\preceq_{(A)}$'s order \mathbb{R}_+^N , we obtain the first inequality. The second inequality requires a little measure theory and Cairoli's maximal inequality [cf. Walsh (1986) for the latter]. We provide a brief but self-contained proof below.

We define one-parameter “filtrations” $\mathcal{F}_1^A, \mathcal{F}_2^A, \dots, \mathcal{F}_N^A$ for each $A \subseteq \Pi$, by insisting that $\mathcal{F}_j^A(t_j)$ is the sigma-field generated by $\{X_j(r); r \geq t_j\}$ if $j \in A$, whereas $\{X_j(r); 0 \leq r \leq t_j\}$ if $j \in A^c$. We add all \mathbb{P}_x -null sets for all $x \in \mathbb{R}^d$ to these without changing our notation. A little thought shows the following: For any $t \in \mathbb{R}_+^N$, given $\mathcal{F}^A(t)$, the sigma-fields $\mathcal{F}_1^A(t_1), \dots, \mathcal{F}_N^A(t_N)$ are conditionally independent under the sigma-finite measure \mathbb{P}_{λ_d} . Consequently, by standard arguments from the theory of Markov random fields, applied to the sigma-finite measure \mathbb{P}_{λ_d} ,

$$\mathbb{E}_{\lambda_d} \{Z | \mathcal{F}^A(t)\} = \mathbb{E}_{\lambda_d} [\dots \mathbb{E}_{\lambda_d} \{Z | \mathcal{F}_N^A(t_N)\} \dots | \mathcal{F}_1^A(t_1)],$$

\mathbb{P}_{λ_d} -a.s. for any nonnegative measurable function Z on Ω [cf. Rozanov (1982), where this sort of result is systematically developed for probability measures]. The

same arguments work for \mathbb{P}_{λ_d} . Consequently, we apply Doob’s maximal inequality, one parameter at a time, to obtain

$$\mathbb{E}_{\lambda_d} \left\{ \sup_{t \in \mathbb{Q}_+^N} |\mathcal{M}_\mu^A f(t)|^2 \right\} \leq 4^N \sup_{t \in \mathbb{R}_+^N} \mathbb{E}_{\lambda_d} \{ |\mathcal{M}_\mu^A f(t)|^2 \},$$

all the time noting that, by applying the method used to prove Kolmogorov’s maximal inequality, one verifies that Doob’s maximal inequality also works for \mathbb{P}_{λ_d} -martingales; we refer to Dellacherie and Meyer [(1978), 40.2, page 34] for the one-parameter discrete setting. One generalizes this development to the multiparameter setting by applying the arguments of Cairoli (1970); cf. Walsh (1986). Thus, Lemma 4.1 concludes our proof. \square

We are ready to begin our proof of the upper bound in Theorem 2.1.

PROOF OF THEOREM 2.1 (Upper bound). Henceforth, we may assume that

$$(4.2) \quad \mathbb{E}\{\lambda_d(X(E))\} > 0,$$

for, otherwise, there is nothing left to prove.

For any $\delta > 0$, define E^δ to be the closed δ -enlargement of E , which is of course a compact set itself. Choose some point $\Delta \notin \mathbb{R}_+^N$, and let T^δ denote any measurable $(\mathbb{Q}_+^N \cap E^\delta) \cup \Delta$ -valued function on Ω , such that $T^\delta \neq \Delta$ if and only if $|X(T^\delta)| \leq \delta$. This can always be done, since the X_j ’s have cadlag paths, and since $\mathcal{B}(0, \delta) = \{x \in \mathbb{R}^d : |x| \leq \delta\}$ has an open interior, where $|\cdot|$ denotes the ℓ^∞ -norm in any Euclidean space. Informally, T^δ is any measurably selected point in E^δ such that $|X(T^\delta)| \leq \delta$, as long as such a point exists. If not, T^δ is set to Δ . We now argue that, for all $\delta > 0$ and all $k > 0$ large, $\mu^{\delta,k} \in \mathcal{P}(E^\delta)$, where

$$\mu^{\delta,k}(\cdot) = \frac{\mathbb{P}_{\lambda_d}\{T^\delta \in \cdot, T^\delta \neq \Delta, |X(0)| \leq k\}}{\mathbb{P}_{\lambda_d}\{T^\delta \neq \Delta, |X(0)| \leq k\}}.$$

Indeed, note that, by Fubini’s theorem,

$$\begin{aligned} \mathbb{P}_{\lambda_d}\{T^\delta \neq \Delta, |X(0)| \leq k\} &= \mathbb{P}_{\lambda_d}\{X(E^\delta) \cap \mathcal{B}(0, \delta) \neq \emptyset, |X(0)| \leq k\} \\ (4.3) \quad &= \int_{[-k, k]^d} \mathbb{P}\{X(E^\delta) \cap \mathcal{B}(x, \delta) \neq \emptyset\} dx \\ &= \mathbb{E}\{\lambda_d((X(E^\delta) \oplus \mathcal{B}(0, \delta)) \cap [-k, k]^d)\}. \end{aligned}$$

In particular, $\mathbb{P}_{\lambda_d}\{T^\delta \neq \Delta, |X(0)| \leq k\}$ is greater than $\mathbb{E}\{\lambda_d(X(E) \cap [-k, k]^d)\} > 0$ for all $k > 0$ large, thanks to (4.2). Thus, once we argue that $\mathbb{P}_{\lambda_d}\{T^\delta \neq \Delta, |X(0)| \leq k\} < +\infty$, this development shows that $\mu^{\delta,k} \in \mathcal{P}(E^\delta)$, as asserted. However,

$$\mathbb{P}_{\lambda_d}\{T^\delta \neq \Delta, |X(0)| \leq k\} \leq \mathbb{P}_{\lambda_d}\{|X(0)| \leq k\} = (2k)^d,$$

which is finite. Thus, we indeed have $\mu^{\delta,k} \in \mathcal{P}(E^\delta)$.

We apply Lemma 4.2 to $\mu = \mu^{\delta,k}$ and $s = T^\delta$ on $\{T^\delta \neq \Delta\}$, and note that on the latter $T^\delta \in \mathbb{Q}_+^N$, so there are no problems with null sets. In this way we obtain the following, where $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ is any measurable function:

$$\sum_{A \subseteq \Pi} \sup_{s \in \mathbb{Q}_+^N} \mathcal{M}_{\mu^{\delta,k}}^A f(s) \geq \int_{\mathbb{R}_+^N} \inf_{x \in \mathbb{R}^d : |x| \leq \delta} P_{t-T^\delta} f(x) \mu^{\delta,k}(dt) \cdot \mathbb{1}_{\{T^\delta \neq \Delta, |X(0)| \leq k\}},$$

\mathbb{P}_{λ_d} -a.s., where the null set is independent of the choice of $\delta > 0$. The special choice of $\mu^{\delta,k}$ yields the following upon squaring and taking \mathbb{P}_{λ_d} -expectations:

$$\begin{aligned} & \mathbb{E}_{\lambda_d} \left\{ \left| \sum_{A \subseteq \Pi} \sup_{s \in \mathbb{Q}_+^N} \mathcal{M}_{\mu^{\delta,k}}^A f(s) \right|^2 \right\} \\ & \geq \int_{\mathbb{R}_+^N} \left[\int_{\mathbb{R}_+^N} \inf_{x \in \mathbb{R}^d : |x| \leq \delta} P_{t-s} f(x) \mu^{\delta,k}(dt) \right]^2 \mu^{\delta,k}(ds) \\ & \quad \times \mathbb{P}_{\lambda_d} \{T^\delta \neq \Delta, |X(0)| \leq k\} \\ & \geq \left[\int_{\mathbb{R}_+^N} \int_{\mathbb{R}_+^N} \inf_{x \in \mathbb{R}^d : |x| \leq \delta} P_{t-s} f(x) \mu^{\delta,k}(dt) \mu^{\delta,k}(ds) \right]^2 \\ & \quad \times \mathbb{P}_{\lambda_d} \{T^\delta \neq \Delta, |X(0)| \leq k\}, \end{aligned}$$

thanks to the Cauchy–Schwarz inequality. Consequently, for all $\delta_0 > 0$, $\delta \in (0, \delta_0)$ and all $k > 0$ large,

$$\begin{aligned} & \mathbb{E}_{\lambda_d} \left\{ \left| \sum_{A \subseteq \Pi} \sup_{s \in \mathbb{Q}_+^N} \mathcal{M}_{\mu^{\delta,k}}^A f(s) \right|^2 \right\} \\ (4.4) \quad & \geq \left[\int_{\mathbb{R}_+^N} \int_{\mathbb{R}_+^N} \inf_{x \in \mathbb{R}^d : |x| \leq \delta_0} P_{t-s} f(x) \mu^{\delta,k}(dt) \mu^{\delta,k}(ds) \right]^2 \\ & \quad \times \mathbb{P}_{\lambda_d} \{T^\delta \neq \Delta, |X(0)| \leq k\}. \end{aligned}$$

Another appeal to the Cauchy–Schwarz inequality reveals the following estimate for the left-hand side of the above:

$$\begin{aligned} (4.5) \quad & \mathbb{E}_{\lambda_d} \left\{ \left| \sum_{A \subseteq \Pi} \sup_{s \in \mathbb{Q}_+^N} \mathcal{M}_{\mu^{\delta,k}}^A f(s) \right|^2 \right\} \leq 2^N \sum_{A \subseteq \Pi} \mathbb{E}_{\lambda_d} \left\{ \left| \sup_{s \in \mathbb{Q}_+^N} \mathcal{M}_{\mu^{\delta,k}}^A f(s) \right|^2 \right\} \\ & \leq 16^N (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{f}(\xi)|^2 \mathcal{Q}_{\mu^{\delta,k}}(\xi) d\xi, \end{aligned}$$

by Lemma 4.2. We now choose a “good” f in both (4.4) and (4.5). Namely, consider $f = f_\varepsilon$ for any $\varepsilon > 0$ such that f is of the form

$$(4.6) \quad f_\varepsilon(x) = (2\pi\varepsilon)^{-d/2} \exp\left(-\frac{\|x\|^2}{2\varepsilon}\right) \quad \forall x \in \mathbb{R}^d.$$

Trivially, $f_\varepsilon \geq 0$, $\int f_\varepsilon d\lambda_d = 1$, $\widehat{f}_\varepsilon(\xi) = \exp(-\frac{1}{2}\varepsilon\|\xi\|^2)$ and both $f_\varepsilon, \widehat{f}_\varepsilon \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$.

We apply (4.4) and (4.5) for this choice of f_ε and wish to take $\delta \rightarrow 0$. Since δ_0 is fixed, the explicit form of $P_t f$ shows that $(x, s, t) \mapsto P_{t-s} f(x)$ is continuous on $\mathcal{B}(0, \delta_0) \times E^\delta \times E^\delta$. Since $E^\delta \downarrow E$ are all compact, and since $\mu^{\delta,k} \in \mathcal{P}(E^\delta)$, Prohorov's theorem, together with the mentioned continuity fact about $\inf_x P_{t-s} f(x)$, implies the existence of a $\mu \in \mathcal{P}(E)$, such that, along some subsequence $\delta' \rightarrow 0$ and $k' \rightarrow \infty$,

$$\begin{aligned} & \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \inf_{|x| \leq \delta_0} P_{t-s} f_\varepsilon(x) \mu^{\delta',k'}(dt) \mu^{\delta',k'}(ds) \\ & \rightarrow \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \inf_{|x| \leq \delta_0} P_{t-s} f_\varepsilon(x) \mu(dt) \mu(ds). \end{aligned}$$

Furthermore, $\mathbb{P}_{\lambda_d}\{T^\delta \neq \Delta, |X(0)| \leq k\} \rightarrow \mathbb{E}\{\lambda_d(\overline{X(E)})\}$, as $k \uparrow \infty$ and then $\delta \downarrow 0$ [cf. (4.3)]. [Here, $\overline{X(E)}$ denotes the closure of $X(E)$.]

The preceding argument, used in conjunction with (4.4) and (4.5) (let $\delta_0 \downarrow 0$), yields

$$\begin{aligned} & 16^N (2\pi)^{-d} \limsup_{\delta' \rightarrow 0, k' \rightarrow \infty} \int_{\mathbb{R}^d} |\widehat{f}_\varepsilon(\xi)|^2 Q_{\mu^{\delta',k'}}(\xi) d\xi \\ & \geq \left[\iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} P_{t-s} f_\varepsilon(0) \mu(dt) \mu(ds) \right]^2 \mathbb{E}\{\lambda_d(X(E))\}. \end{aligned}$$

Now, $0 \leq Q_{\mu^{\delta',k'}}(\xi) \leq 1$ for all $\xi \in \mathbb{R}^d$ [cf. (3.7)], and $\widehat{f}_\varepsilon \in L^2(\mathbb{R}^d)$. Finally,

$$\lim_{\delta' \rightarrow 0, k' \rightarrow \infty} Q_{\mu^{\delta',k'}} = Q_\mu,$$

pointwise [cf. (3.6)]. Thus, by the Lebesgue dominated convergence theorem,

$$\begin{aligned} & 16^N (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{f}_\varepsilon(\xi)|^2 Q_\mu(\xi) d\xi \\ & \geq \left[\iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} P_{t-s} f_\varepsilon(0) \mu(dt) \mu(ds) \right]^2 \mathbb{E}\{\lambda_d(X(E))\} \\ & = (2\pi)^{-2d} \left[\int_{\mathbb{R}^d} \widehat{f}_\varepsilon(\xi) Q_\mu(\xi) d\xi \right]^2 \mathbb{E}\{\lambda_d(X(E))\}, \end{aligned}$$

thanks to Lemma 3.1. Our choice of f_ε guarantees that, for all $\xi \in \mathbb{R}^d$, $\widehat{f}_\varepsilon(\xi) \geq |\widehat{f}_\varepsilon(\xi)|^2$. Thus, we use the square integrability of f_ε and the positivity and boundedness of Q_μ , once more, to obtain

$$\mathbb{E}\{\lambda_d(X(E))\} \leq 16^N \left[(2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{f}_\varepsilon(\xi)|^2 Q_\mu(\xi) d\xi \right]^{-1},$$

since the right-hand side is obviously not zero [cf. (3.6) and (3.7)]. Let $\varepsilon \downarrow 0$ and use the Lebesgue monotone convergence theorem to see that

$$\begin{aligned} \mathbb{E}\{\lambda_d(X(E))\} &\leq 16^N \left[(2\pi)^{-d} \int_{\mathbb{R}^d} Q_\mu(\xi) d\xi \right]^{-1} \\ &= 16^N \left[(2\pi)^{-d} \mathbb{E}\left\{ \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2 \right\} \right]^{-1}, \end{aligned}$$

by Lemma 2.4. This concludes our proof. \square

5. Proofs of Theorems 1.1, 1.3 and 1.5. Theorem 1.3 follows from Theorem 1.1 by invoking the very argument that lead to Corollary 2.3. Hence, we only concentrate on proving Theorems 1.1 and 1.5.

We divide the proofs of Theorems 1.1 and 1.5 into three parts:

1. The easy half of Theorem 1.1, that is,

$$\int_{\mathbb{R}^d} \prod_{j=1}^N \operatorname{Re} \left(\frac{1}{1 + \Psi_j(\xi)} \right) d\xi < +\infty \implies \mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} > 0.$$

Of course, this statement is a special case of part 3 below. We give a simple and direct proof using Theorem 2.1.

2. The hard half of Theorem 1.5; that is, for any fixed compact set $F \subset \mathbb{R}^d$,

$$\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N) \oplus F)\} > 0 \implies \exists \mu \in \mathcal{P}(F) \text{ such that } \mathcal{E}_\Psi(\mu) < \infty.$$

The hard half of Theorem 1.1 follows from this and (1.4) upon selecting $F = \{0\}$.

3. The easy half of Theorem 1.5; that is,

$$F \text{ carries a finite measure of finite energy} \implies \mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N) \oplus F)\} > 0.$$

For simplicity, we use the following suggestive notation: For all finite measures μ on \mathbb{R}_+^N , define

$$\|\mu\|_e^2 = (2\pi)^{-d} \mathbb{E}\left\{ \|\widehat{\mathbb{O}}_\mu\|_{L^2(\mathbb{R}^d)}^2 \right\}.$$

We may refer to $\|\mu\|_e$ as the *energy norm* of μ , although strictly speaking it is only a seminorm as the following shows.

LEMMA 5.1. *For any two finite measures μ and ν on \mathbb{R}^d ,*

$$\|\mu + \nu\|_e \leq \|\mu\|_e + \|\nu\|_e.$$

PROOF. Since $\mu \mapsto \mathbb{O}_\mu$ is linear, so is $\mu \mapsto \widehat{\mathbb{O}}_\mu$. The lemma follows from Minkowski's inequality. \square

Throughout, we define the *one-killing measure* $\kappa \in \mathcal{P}(\mathbb{R}_+^N)$ as

$$(5.1) \quad \kappa(ds) = \exp\left(-\sum_{j=1}^N s_j\right) ds \quad \forall s \in \mathbb{R}_+^N.$$

Recalling (2.1), we note the *killed occupation measure* is defined by

$$\mathbb{O}_\kappa(f) = \int_{\mathbb{R}_+^N} f(X(s)) \exp\left(-\sum_{j=1}^N s_j\right) ds.$$

Note that \mathbb{O}_κ is a random probability measure carried by $X(\mathbb{R}_+^N)$.

The relevance of the killing measure κ to the proofs of Theorems 1.1 and 1.5 is given by the following lemma.

LEMMA 5.2. *Let κ be the killing measure defined above. Then the energy norm of $\kappa \in \mathcal{P}(\mathbb{R}_+^N)$ is described by*

$$\|\kappa\|_e^2 = (2\pi)^{-d} \int_{\mathbb{R}^d} \prod_{j=1}^N \operatorname{Re}\left(\frac{1}{1 + \Psi_j(\xi)}\right) d\xi.$$

PROOF. By Lemma 2.4,

$$\begin{aligned} \|\kappa\|_e^2 &= (2\pi)^{-d} \int_{\mathbb{R}^d} \prod_{j=1}^N \left[\int_0^\infty \int_0^\infty e^{-|s-t|\Psi_j(\operatorname{sgn}(s-t)\xi)} e^{-s-t} ds dt \right] d\xi \\ &= (2\pi)^{-d} \int_{\mathbb{R}^d} \prod_{j=1}^N \left[\int_0^\infty \int_s^\infty (\dots) dt ds + \int_0^\infty \int_t^\infty (\dots) ds dt \right] d\xi \\ &= (2\pi)^{-d} \int_{\mathbb{R}^d} \prod_{j=1}^N \left[\frac{1/2}{1 + \Psi_j(\xi)} + \frac{1/2}{1 + \overline{\Psi_j(\xi)}} \right] d\xi \\ &= (2\pi)^{-d} \int_{\mathbb{R}^d} \prod_{j=1}^N \operatorname{Re}\left(\frac{1}{1 + \Psi_j(\xi)}\right) d\xi, \end{aligned}$$

since $\{1 + z\}^{-1} + \{1 + \bar{z}\}^{-1} = 2 \operatorname{Re}\{1 + z\}^{-1}$ ($z \in \mathbb{C}$). \square

Lemma 5.2 suffices for our proof of the easy half of Theorem 1.1.

PROOF OF THEOREM 1.1 (Easy half). The lower bound in Theorem 2.1 shows that

$$(5.2) \quad \mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} \geq \|\kappa\|_e^{-2}.$$

Thus, Lemma 5.2 shows that the integrability of $\prod_{j=1}^N \operatorname{Re}\{1 + \Psi_j\}^{-1}$ guarantees the positivity of $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\}$. This completes part 1 of the proof. \square

We start working toward proving the hard half of Theorem 1.5. We begin with some prefatory results.

Recalling our definition of energy from (1.4), and the function Q_κ from (3.6), we have the following.

LEMMA 5.3. *For all $\xi \in \mathbb{R}^d$,*

$$Q_\kappa(\xi) = \prod_{j=1}^N \operatorname{Re}\left(\frac{1}{1 + \Psi_j(\xi)}\right).$$

Consequently, whenever $f: \mathbb{R}^d \rightarrow \mathbb{R}$ and its Fourier transform are both in $L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$,

$$\mathbb{E}_{\lambda_d}\{|\mathbb{O}_\kappa(f)|^2\} = \mathcal{E}_\Psi(f).$$

PROOF. In light of Lemma 3.5 and our definition of energy (1.4), it suffices to compute Q_κ as given. On the other hand,

$$\begin{aligned} Q_\kappa(\xi) &= \prod_{j=1}^N \int_0^\infty \int_0^\infty e^{-s-t-|s-t|\Psi_j(\operatorname{sgn}(s-t)\xi)} ds dt \\ &= \prod_{j=1}^N \left[\int_0^\infty \int_t^\infty (\dots) ds dt + \int_0^\infty \int_s^\infty (\dots) dt ds \right] \\ &= \prod_{j=1}^N \operatorname{Re}\left(\frac{1}{1 + \Psi_j(\xi)}\right). \end{aligned}$$

The few remaining details in the above are the same as those in the proof of Lemma 5.2. \square

Recalling (4.1), we are interested in $\mathcal{M}_\kappa^\Pi f$, where we also recall that $\Pi = \{1, \dots, N\}$. The operator U_+ below was defined in (3.3).

LEMMA 5.4. *If $f: \mathbb{R}^d \rightarrow \mathbb{R}_+$ is in $L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$, then*

$$(5.3) \quad \mathbb{E}_{\lambda_d} \left\{ \sup_{s \in \mathbb{Q}_+^N} |\mathcal{M}_\kappa^\Pi f(s)|^2 \right\} \leq 4^N \mathcal{E}_\Psi(f).$$

Furthermore, for any $r > 0$, for any $f \in L^1(\mathbb{R}^d)$ whose Fourier transform is also in $L^1(\mathbb{R}^d)$ and for all $s \in (0, r)^N$, the following holds \mathbb{P}_{λ_d} -a.s.:

$$(5.4) \quad \mathcal{M}_\kappa^\Pi f(s) \geq e^{-Nr} U_+ f(X(s)).$$

PROOF. Equation (5.3) is a consequence of Lemmas 4.2 and 5.3 albeit in slightly different notation.

To prove (5.4), we proceed as in our proof of Lemma 4.2, but adapt the argument to the present setting. Since $f \geq 0$, the same reasoning as in the latter lemma gives the following for all $s \in (0, r)^N$:

$$\begin{aligned} \mathcal{M}_k^\Pi f(s) &\geq \mathbb{E}_{\lambda_d} \left\{ \int_{t \succ_{(\Pi)} s} f(X(t)) \exp\left(-\sum_{j=1}^N t_j\right) dt \middle| \mathcal{F}^\Pi(s) \right\} \\ &= \int_{t \succ_{(\Pi)} s} P_{t-s} f(X(s)) \exp\left(-\sum_{j=1}^N t_j\right) dt, \end{aligned}$$

\mathbb{P}_{λ_d} -a.s. Now we move in a somewhat new direction by noticing that, since $s \in (0, r)^N$,

$$\begin{aligned} \mathcal{M}_k^\Pi f(s) &\geq e^{-Nr} \int_{t \succ_{(\Pi)} s} P_{t-s} f(X(s)) \exp\left(-\sum_{j=1}^N (t_j - s_j)\right) dt \\ &\geq e^{-Nr} \int_{\mathbb{R}_+^N} P_t f(X(s)) \exp\left(-\sum_{j=1}^N t_j\right) dt \\ &= e^{-Nr} U_+ f(X(s)), \end{aligned}$$

by (3.3). Thus, our lemma follows. \square

Henceforth, we define the *capacity* of a compact set $F \subset \mathbb{R}^d$ by

$$(5.5) \quad \mathcal{C}_\Psi(F) = \left[\inf_{\mu \in \mathcal{P}(F)} \mathcal{E}_\Psi(\mu) \right]^{-1},$$

where, we recall, $\mathcal{P}(F)$ denotes the collection of all probability measures on F .

Our proof of the hard half of Theorem 1.5 is based on the following.

LEMMA 5.5. *Suppose X is an additive Lévy process in \mathbb{R}^d that satisfies condition (1.3), and that $\int_{\mathbb{R}^d} \prod_{j=1}^N |1 + \Psi_j(\xi)|^{-1} d\xi < +\infty$, where $\Psi = (\Psi_1, \dots, \Psi_N)$ denotes the Lévy exponent of X . Then, for all compact sets $F \subset \mathbb{R}^d$, and for all $r > 0$,*

$$\mathbb{E}\{\lambda_d(X([0, r]^N) \oplus F)\} \leq \vartheta^{-2} (4e^{2r})^N \cdot \mathcal{C}_\Psi(F),$$

where $\vartheta > 0$ is the constant in condition (1.3).

Before proving it, we appeal to Lemma 5.5 to conclude part 2 of our proof, that is, the hard half of Theorem 1.5. Clearly, the following will suffice.

PROPOSITION 5.6. *Suppose X is an additive Lévy process in \mathbb{R}^d that satisfies condition (1.3). Let $\Psi = (\Psi_1, \dots, \Psi_N)$ denote the Lévy exponent of X . Then, for all compact sets $F \subset \mathbb{R}^d$ and for all $r > 0$,*

$$\mathbb{E}\{\lambda_d(X([0, r]^N) \oplus F)\} \leq \vartheta^{-2}(4e^{2r})^{N+\lfloor d/2 \rfloor+1} \cdot \mathcal{C}_\Psi(F).$$

PROOF. Let us bring in $M = \lfloor \frac{d}{2} \rfloor + 1$ continuous Brownian motions in \mathbb{R}^d , B_1, \dots, B_M , all totally independent from one another, as well as X [under \mathbb{P}]. Having done so, for any $\delta > 0$, we can define an $(N + M)$ -parameter process \mathcal{X}_δ in \mathbb{R}^d by

$$\mathcal{X}_\delta = X_1 \oplus \dots \oplus X_N \oplus \sqrt{2\delta}B_1 \oplus \dots \oplus \sqrt{2\delta}B_M.$$

To be concrete, $\mathcal{X}_\delta(t) = \sum_{j=1}^N X_j(t_j) + \sqrt{2\delta} \sum_{j=1}^M B_j(t_{j+N})$, for all $t \in \mathbb{R}_+^{N+M}$.

Define

$$\Phi_j(\xi) = \begin{cases} \Psi_j(\xi), & \text{if } j = 1, \dots, N, \\ \delta \|\xi\|^2, & \text{if } j = N + 1, \dots, N + M, \end{cases}$$

where $\|x\|$ is the ℓ^2 -norm of $x \in \mathbb{R}^d$. Then \mathcal{X}_δ is an $(N + M)$ -parameter additive Lévy process in \mathbb{R}^d whose Lévy exponent is $\Phi = (\Phi_1, \dots, \Phi_{N+M})$. Furthermore,

$$(5.6) \quad \int_{\mathbb{R}^d} \prod_{j=1}^{N+M} |1 + \Phi_j(\xi)|^{-1} d\xi \leq \int_{\mathbb{R}^d} \{1 + \delta \|\xi\|^2\}^{-M} d\xi < +\infty,$$

since $M > \frac{d}{2}$. Thus, we can apply Lemma 5.5 to the process \mathcal{X}_δ and obtain

$$(5.7) \quad \mathbb{E}\{\lambda_d(\mathcal{X}_\delta([0, r]^{N+M}) \oplus F)\} \leq \vartheta^{-2}(4e^{2r})^{N+M} \cdot \mathcal{C}_\Phi(F).$$

Of course, by (5.6), the above capacity is strictly positive.

Now, consider a sequence of probability measures, μ_1, μ_2, \dots , all on F , such that

$$\lim_{n \rightarrow \infty} \mathcal{E}_\Phi(\mu_n) = [\mathcal{C}_\Phi(F)]^{-1}.$$

Without loss of generality, we can assume that all these energies are finite and, by tightness, extract a subsequence n' and a probability measure μ_∞ on F such that $\mu_{n'}$ converges weakly to μ_∞ . Thanks to (5.6) and to the continuity of the Φ_j 's, we see that $\lim_{n'} \mathcal{E}_\Phi(\mu_{n'}) = \mathcal{E}_\Phi(\mu_\infty)$. Equivalently, we have found a probability measure μ_∞ on F whose Φ -energy is the reciprocal of the Φ -capacity of F . Changing the notation to allow for the dependence of μ_∞ on the parameter δ , we see that there exists a probability measure ν_δ on F such that it has finite Φ -energy, and

$$\mathbb{E}\{\lambda_d(\mathcal{X}_\delta([0, r]^{N+M}) \oplus F)\} \leq \frac{\vartheta^{-2}(4e^{2r})^{N+M}}{\mathcal{E}_\Phi(\nu_\delta)}.$$

This holds for all $\delta > 0$. Now, fix an arbitrarily small $\delta_0 > 0$, and for all $\delta \in (0, \delta_0)$ deduce the cruder bound:

$$\begin{aligned} & \mathbb{E}\{\lambda_d(\mathcal{X}_\delta([0, r]^{N+M}) \oplus F)\} \\ & \leq \frac{\vartheta^{-2}(4e^{2r})^{N+M}}{\int_{\mathbb{R}^d} |\widehat{v}_\delta(\xi)|^2 \prod_{j=1}^N \operatorname{Re}\{1 + \Psi_j(\xi)\}^{-1} \cdot \{1 + \delta_0 \|\xi\|^2\}^{-M} d\xi}. \end{aligned}$$

By considering further subnets of δ , and by appealing to (5.6) once more, we can infer the existence of a probability measure ν_0 on F such that

$$\begin{aligned} & \liminf_{\delta \rightarrow 0} \mathbb{E}\{\lambda_d(\mathcal{X}_\delta([0, r]^{N+M}) \oplus F)\} \\ & \leq \frac{\vartheta^{-2}(4e^{2r})^{N+M}}{\int_{\mathbb{R}^d} |\widehat{\nu}_0(\xi)|^2 \prod_{j=1}^N \operatorname{Re}\{1 + \Psi_j(\xi)\}^{-1} \cdot \{1 + \delta_0 \|\xi\|^2\}^{-M} d\xi}. \end{aligned}$$

On the other hand,

$$\mathcal{X}_\delta([0, r]^{N+M}) = X([0, r]^N) \oplus \sqrt{2\delta}B([0, r]^M),$$

where $B = B_1 \oplus \dots \oplus B_M$. By compactness, as $\delta \downarrow 0$, this random set converges downward to $\overline{X([0, r]^N)}$, the closure of $X([0, r]^N)$. Consequently, by the monotone convergence theorem of Lebesgue,

$$\begin{aligned} & \mathbb{E}\{\lambda_d(X([0, r]^N) \oplus F)\} \\ & \leq \frac{\vartheta^{-2}(4e^{2r})^{N+M}}{\int_{\mathbb{R}^d} |\widehat{\nu}_0(\xi)|^2 \prod_{j=1}^N \operatorname{Re}\{1 + \Psi_j(\xi)\}^{-1} \cdot \{1 + \delta_0 \|\xi\|^2\}^{-M} d\xi}, \end{aligned}$$

for all $\delta_0 > 0$. Let $\delta_0 \downarrow 0$, and apply Lebesgue monotone convergence one more time to finish. \square

It remains to present our proof of Lemma 5.5.

PROOF OF LEMMA 5.5. It follows from (1.4) that $\mathcal{C}_\Psi(-F) = \mathcal{C}_\Psi(F)$ for any compact set F , where $-F = \{-a; a \in F\}$. Hence, we can reduce our problem to showing that

$$\mathbb{E}\{\lambda_d(F \ominus X([0, r]^N))\} \leq \vartheta^{-2}(4e^{2r})^N \mathcal{C}_\Psi(F),$$

where $A \ominus B = \{a - b; a \in A, b \in B\}$.

Let F^ε denote the closed ε -enlargement of F . The integrability condition of the statement of the lemma, the continuity of the Ψ_j 's and the Lebesgue dominated convergence theorem together show that if $\varepsilon > 0$, then $\lim_{\varepsilon \rightarrow 0} \mathcal{C}_\Psi(F^\varepsilon) = \mathcal{C}_\Psi(F)$. (This involves a tightness argument that we have already utilized while proving Proposition 5.6.) Hence, it suffices to show that

$$(5.8) \quad \mathbb{E}\{\lambda_d(F \ominus X([0, r]^N))\} \leq \vartheta^{-2}(4e^{2r})^N \mathcal{C}_\Psi(F^\varepsilon).$$

The above holds trivially unless the left-hand side is strictly positive, which we will assume henceforth. We observe that, by Fubini’s theorem,

$$\begin{aligned}
 \mathbb{P}_{\lambda_d}\{X([0, r]^N) \cap F \neq \emptyset\} &= \int_{\mathbb{R}^d} \mathbb{P}\{(x \oplus X([0, r]^N)) \cap F \neq \emptyset\} dx \\
 (5.9) \qquad \qquad \qquad &= \int_{\mathbb{R}^d} \mathbb{P}\{x \in F \ominus X([0, r]^N)\} dx \\
 &= \mathbb{E}\{\lambda_d(F \ominus X([0, r]^N))\}.
 \end{aligned}$$

Thus, the above assumption is equivalent to assuming that

$$(5.10) \qquad \qquad \mathbb{P}_{\lambda_d}\{X([0, r]^N) \cap F \neq \emptyset\} > 0.$$

Next, we add a cemetery point $\Delta \notin \mathbb{R}_+^N$ to \mathbb{R}_+^N , and consider any $\mathbb{Q}_+^N \cup \{\Delta\}$ -valued random variable T_ε such that the following hold:

- (i) $T_\varepsilon = \Delta$ if and only if $X([0, r]^N) \cap F^\varepsilon = \emptyset$;
- (ii) on the event $\{X([0, r]^N) \cap F^\varepsilon \neq \emptyset\}$, $X(T_\varepsilon) \in F^\varepsilon$.

We remark that since F^ε has an open interior, and since X_i ’s are cadlag, we can always choose $T_\varepsilon \in \mathbb{Q}_+^N \cup \{\Delta\}$ (as opposed to $\mathbb{R}_+^N \cup \{\Delta\}$). Consider

$$\begin{aligned}
 \mu_{\varepsilon,k}(\cdot) &= \frac{\mathbb{P}_{\lambda_d}\{X(T_\varepsilon) \in \cdot, T_\varepsilon \neq \Delta, |X(0)| \leq k\}}{\mathbb{P}_{\lambda_d}\{T_\varepsilon \neq \Delta, |X(0)| \leq k\}}, \\
 \varphi_\eta(x) &= (2\pi\eta^2)^{-d/2} \exp\left(-\frac{\|x\|^2}{2\eta^2}\right),
 \end{aligned}$$

where $\eta, k > 0$ and $x \in \mathbb{R}^d$. Owing to (5.10), $\mu_{\varepsilon,k}$ is a probability measure on F^ε for all $\varepsilon > 0$ and $k > 0$ large. We can smooth $\mu_{\varepsilon,k}$ by convoluting it with φ_η :

$$f_{\varepsilon,k;\eta} = \mu_{\varepsilon,k} \star \varphi_\eta.$$

The function $f_{\varepsilon,k;\eta}$ has the following nice properties that are simple to check:

- (i) $f_{\varepsilon,k;\eta} \geq 0$ is bounded; and
- (ii) both $f_{\varepsilon,k;\eta}$ and $\widehat{f_{\varepsilon,k;\eta}} \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$. Thus, we can apply Lemma 5.4 [(5.4)] to obtain

$$\sup_{t \in [0, r]^N} \mathcal{M}_\kappa^\Pi f_{\varepsilon,k;\eta}(t) \geq e^{-Nr} U_+ f_{\varepsilon,k;\eta}(X(T_\varepsilon)) \cdot \mathbb{1}_{\{T_\varepsilon \neq \Delta, |X(0)| \leq k\}},$$

\mathbb{P}_{λ_d} -a.s. (There are no problems with null sets, since on $\{T_\varepsilon \neq \Delta\}$, $T_\varepsilon \in \mathbb{Q}_+^N$.) We square this and take \mathbb{E}_{λ_d} -expectations to obtain the following as a consequence of

Lemma 5.4 [(5.3)]:

$$\begin{aligned}
 & 4^N \mathcal{E}_\Psi(f_{\varepsilon,k;\eta}) \\
 & \geq e^{-2Nr} \mathbb{E}_{\lambda_d} \left\{ |U_+ f_{\varepsilon,k;\eta}(X(T_\varepsilon))|^2 \mathbb{1}_{\{T_\varepsilon \neq \Delta, |X(0)| \leq k\}} \right\} \\
 (5.11) \quad & = e^{-2Nr} \int_{\mathbb{R}^d} |U_+ f_{\varepsilon,k;\eta}(x)|^2 \mu_{\varepsilon,k}(dx) \cdot \mathbb{P}_{\lambda_d} \{T_\varepsilon \neq \Delta, |X(0)| \leq k\} \\
 & \geq e^{-2Nr} \left| \int_{\mathbb{R}^d} U_+ f_{\varepsilon,k;\eta}(x) \mu_{\varepsilon,k}(dx) \right|^2 \cdot \mathbb{P}_{\lambda_d} \{T_\varepsilon \neq \Delta, |X(0)| \leq k\},
 \end{aligned}$$

thanks to the Cauchy–Schwarz inequality. We can apply Lemma 3.1 to see that

$$\begin{aligned}
 & \int_{\mathbb{R}^d} U_+ f_{\varepsilon,k;\eta}(x) \mu_{\varepsilon,k}(dx) \\
 & = (2\pi)^{-d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \widehat{f_{\varepsilon,k;\eta}}(-\xi) \prod_{j=1}^N \frac{1}{1 + \Psi_j(\xi)} d\xi \mu_{\varepsilon,k}(dx) \\
 & = (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{\mu_{\varepsilon,k}}(\xi)|^2 \widehat{\varphi}_\eta(-\xi) \operatorname{Re} \left(\prod_{j=1}^N \frac{1}{1 + \Psi_j(\xi)} \right) d\xi \\
 & \geq \vartheta (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{\mu_{\varepsilon,k}}(\xi)|^2 \exp\left(-\frac{\|x\|^2}{2\eta^2}\right) \prod_{j=1}^N \operatorname{Re} \left(\frac{1}{1 + \Psi_j(\xi)} \right) d\xi \\
 & \xrightarrow{\eta \rightarrow 0} \vartheta \mathcal{E}_\Psi(\mu_{\varepsilon,k}).
 \end{aligned}$$

In the above, the second equality follows from the fact that $|\widehat{\mu_{\varepsilon,k}}(\xi)|^2 \widehat{\varphi}_\eta(-\xi) \geq 0$, and the inequality follows from condition (1.3). On the other hand, by (5.9),

$$\begin{aligned}
 & \lim_{k \rightarrow \infty} \mathbb{P}_{\lambda_d} \{T_\varepsilon \neq \Delta, |X(0)| \leq k\} \\
 & = \mathbb{P}_{\lambda_d} \{T_\varepsilon \neq \Delta\} \\
 & = \mathbb{E} \{ \lambda_d(F^\varepsilon \ominus X([0, r]^N)) \} \\
 & \geq \mathbb{E} \{ \lambda_d(F \ominus X([0, r]^N)) \},
 \end{aligned}$$

since $F \subset F^\varepsilon$. Finally, the integrability condition of our lemma allows us to take limits and conclude that $\lim_{\eta \rightarrow 0} \mathcal{E}_\Psi(f_{\varepsilon,k;\eta}) = \mathcal{E}_\Psi(\mu_{\varepsilon,k})$. Thus, (5.11) implies (5.8) after letting $k \uparrow \infty$, from which our lemma follows. \square

Next, we proceed to part 3 of our proof. The following proposition verifies the easy half of Theorem 1.5. Namely, if F carries a finite measure with finite energy, then the Lebesgue measure of $X(\mathbb{R}_+^N) \oplus F$ has a positive expectation.

PROPOSITION 5.7. *Suppose X is an additive Lévy process in \mathbb{R}^d with Lévy exponent $\Psi = (\Psi_1, \dots, \Psi_N)$. If $F \subset \mathbb{R}^d$ is a compact set and $\mathcal{C}_\Psi(F) > 0$, then for any $r > 0$,*

$$\mathbb{E}\{\lambda_d(X([0, r]^N) \oplus F)\} > 0.$$

PROOF. Since $\mathcal{C}_\Psi(-F) = \mathcal{C}_\Psi(F)$, as in the proof of Lemma 5.5 we only need to show that if $\mathcal{C}_\Psi(F) > 0$, then for any $r > 0$,

$$(5.12) \quad \mathbb{E}\{\lambda_d(X([0, r]^N) \ominus F)\} > 0.$$

We note that, whenever $\mathbb{E}\{\lambda_d(X([0, r]^N))\} > 0$, then $\mathbb{E}\{\lambda_d(X([0, r]^N) \ominus F)\} > 0$ for all compact sets $F \subset \mathbb{R}^d$, and the proposition holds trivially. Moreover, if there exist some $n \leq N - 1$, distinct $i_1, \dots, i_n \in \{1, \dots, N\}$ and a compact set $F \subset \mathbb{R}^d$ such that

$$\mathbb{E}\{\lambda_d(X_{i_1 \dots i_n}([0, r]^n) \ominus F)\} > 0,$$

where $X_{i_1 \dots i_n} = X_{i_1} \oplus \dots \oplus X_{i_n}$ is an n -parameter additive Lévy process, then inequality (5.12) also holds. Hence, without loss of generality, we can and will assume that

$$(5.13) \quad \mathbb{E}\{\lambda_d(X_{i_1 \dots i_n}([0, r]^n) \ominus F)\} = 0$$

for all $n \leq N - 1$ and distinct $i_1, \dots, i_n \in \{1, \dots, N\}$.

For any $\varepsilon > 0$, define

$$\varphi_\varepsilon(x) = (2\varepsilon)^{-d} \mathbb{1}_{\{|x| \leq \varepsilon\}}, \quad x \in \mathbb{R}^d,$$

where $|x|$ is the ℓ^∞ -norm of $x \in \mathbb{R}^d$. Then, whenever μ is a probability measure on F , $\mu_\varepsilon := \mu \star \varphi_\varepsilon$ is a probability measure on F^ε , where \star denotes convolution, and F^ε is the closed ε -enlargement of F in the ℓ^∞ -norm. To maintain some notational simplicity, we write μ_ε both for the measure and its density with respect to Lebesgue measure λ_d .

Since the \mathbb{P}_{λ_d} -distribution of $X(t)$ is λ_d for all $t \in \mathbb{R}_+^N$,

$$(5.14) \quad \mathbb{E}_{\lambda_d} \left\{ \int_{[0, r]^N} \exp\left(-\sum_{j=1}^N s_j\right) \mu_\varepsilon(X(s)) ds \right\} = (1 - e^{-r})^N.$$

On the other hand, by Lemma 5.3 and the definition of energy (1.4),

$$(5.15) \quad \mathbb{E}_{\lambda_d} \left\{ \left[\int_{[0, r]^N} \exp\left(-\sum_{j=1}^N s_j\right) \mu_\varepsilon(X(s)) ds \right]^2 \right\} \leq \mathcal{E}_\Psi(\mu_\varepsilon) \leq \mathcal{E}_\Psi(\mu),$$

since $|\widehat{\varphi_\varepsilon}(\xi)| \leq 1$. Recall the Paley–Zygmund inequality: For any measure ν on the underlying measure space, and for any nonnegative $g \in L^2(\nu) \cap L^1(\nu)$,

$$\nu\{g > 0\} \geq \frac{\|g\|_{L^1(\nu)}^2}{\|g\|_{L^2(\nu)}^2}$$

[cf. Kahane (1985), page 8]. We apply this with $\nu = \mathbb{P}_{\lambda_d}$ and

$$g(\omega) = \int_{[0,r]^N} \exp\left(-\sum_{j=1}^N s_j\right) \mu_\varepsilon(X(s))(\omega) ds.$$

Thanks to (5.14) and (5.15), we have $\|g\|_{L^1(\nu)} = (1 - e^{-r})^N$ and $\|g\|_{L^2(\nu)}^2 \leq \mathcal{E}_\Psi(\mu)$. Therefore,

$$\mathbb{P}_{\lambda_d}\{X([0, r]^N) \cap F^\varepsilon \neq \emptyset\} \geq \mathbb{P}_{\lambda_d}\{g > 0\} \geq (1 - e^{-r})^{2N} [\mathcal{E}_\Psi(\mu)]^{-1}.$$

It follows from (5.9) that

$$\mathbb{P}_{\lambda_d}\{X([0, r]^N) \cap F^\varepsilon \neq \emptyset\} = \mathbb{E}\{\lambda_d(X([0, r]^N) \ominus F^\varepsilon)\}.$$

Thus, we can let $\varepsilon \downarrow 0$ to obtain

$$(5.16) \quad \mathbb{E}\{\lambda_d(\overline{X([0, r]^N)} \ominus F)\} \geq (1 - e^{-r})^{2N} [\mathcal{E}_\Psi(\mu)]^{-1}$$

for all probability measures μ on F that have finite energy. Since each X_j has only a countable number of jumps, the assumption (5.13) implies that

$$\lambda_d\left(\left(\overline{X([0, r]^N)} \setminus X([0, r]^N)\right) \ominus F\right) = 0, \quad \mathbb{P}\text{-a.s.}$$

Therefore, (5.16) becomes

$$\mathbb{E}\{\lambda_d(X([0, r]^N) \ominus F)\} \geq (1 - e^{-r})^{2N} [\mathcal{E}_\Psi(\mu)]^{-1}$$

for all probability measures μ on F that have finite energy. Defining $1 \div 0 = \infty$ as we have, we can optimize over all probability measures μ on F to deduce that

$$\mathbb{E}\{\lambda_d(X([0, r]^N) \ominus F)\} \geq (1 - e^{-r})^{2N} \mathcal{C}_\Psi(F).$$

This proves (5.12), and our proposition follows. \square

6. Convolution-based energies. This is a brief section on connections between the energy forms of the Introduction and the notion of mutual energy based on convolutions. Some of this material is classical and can be found in standard references such as Carleson (1983) and Kahane (1985).

Any locally integrable function $K : \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{R}_+$ defines a mutual energy on the space of all measures crossed itself. To be precise, the K -mutual energy of measures μ and ν is defined by

$$(6.1) \quad (\mu, \nu)_K = \frac{1}{2} \iint K(a - b) \mu(da) \nu(db) + \frac{1}{2} \iint K(b - a) \mu(da) \nu(db).$$

This is clearly a symmetric form, that is, $(\mu, \nu)_K = (\nu, \mu)_K$. It also induces a capacity \mathbf{C}_K on subsets of \mathbb{R}^d :

$$\mathbf{C}_K(F) = \left[\inf_{\mu \in \mathcal{P}(F)} (\mu, \mu)_K \right]^{-1},$$

where $\mathcal{P}(F)$ is the collection of all probability measures that are carried by F .

We say that K is the 1-potential density of an additive Lévy process $X = \{X(t); t \in \mathbb{R}_+^N\}$ in \mathbb{R}^d if, for all $f: \mathbb{R}^d \rightarrow \mathbb{R}_+$,

$$\mathbb{E} \left\{ \int_{\mathbb{R}_+^N} \exp \left(- \sum_{j=1}^N s_j \right) f(X(s)) ds \right\} = \int_{\mathbb{R}^d} K(a) f(a) da.$$

It is easy to see that if the 1-potential density $K(a)$ exists, then $\int_{\mathbb{R}^d} K(a) da = 1$ and $K(a) > 0$ for almost every $a \in X(\mathbb{R}_+^N)$. A sufficient condition for the existence of the one-potential density is that $X(s)$ has a density function $p_s(a)$ for all $s \in (0, \infty)^N$. In this case,

$$(6.2) \quad K(a) = \int_{\mathbb{R}_+^N} \exp \left(- \sum_{j=1}^N s_j \right) p_s(a) ds.$$

See Hawkes [(1979), Lemma 2.1] for a necessary and sufficient condition for the existence of a one-potential density.

LEMMA 6.1. *Let X be an additive Lévy process in \mathbb{R}^d with Lévy exponent Ψ and 1-potential density K , and suppose condition (1.3) holds. Then, for all finite measures μ on \mathbb{R}^d ,*

$$2^{-N}(\mu, \mu)_K \leq \mathcal{E}_\Psi(\mu) \leq \vartheta^{-1}(\mu, \mu)_K.$$

Furthermore, if (1.7) holds, then $\mathcal{E}_\Psi(\mu) = (\mu, \mu)_K$.

PROOF. Define $K_\star(a) = \frac{1}{2}[K(a) + K(-a)]$ to be the symmetrization of K . Then

$$(6.3) \quad \begin{aligned} \widehat{K}_\star(\xi) &= \frac{1}{2} \left[\prod_{j=1}^N \frac{1}{1 + \Psi_j(\xi)} + \prod_{j=1}^N \frac{1}{1 + \overline{\Psi_j(\xi)}} \right] \\ &= \operatorname{Re} \left(\prod_{j=1}^N \frac{1}{1 + \Psi_j(\xi)} \right). \end{aligned}$$

We note that \widehat{K}_\star is a real function and, under condition (1.3), it is also nonnegative. Hence, by Fubini's theorem,

$$(6.4) \quad \begin{aligned} (\mu, \mu)_K &= \iint_{\mathbb{R}^d \times \mathbb{R}^d} K_\star(a - b) \mu(da) \mu(db) \\ &= (2\pi)^{-d} \iint \int e^{-i\xi \cdot (a-b)} \widehat{K}_\star(\xi) d\xi \mu(da) \mu(db) \\ &= (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^2 \widehat{K}_\star(\xi) d\xi. \end{aligned}$$

Together, this and (6.3) imply the second portion of the lemma, as well as the

asserted upper bound on $\mathcal{E}_\Psi(\mu)$. It remains to verify the corresponding lower bound for $\mathcal{E}_\Psi(\mu)$.

Note that if $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ is measurable, $U_+ f \leq Uf$, pointwise [cf. (3.2) and (3.3)]. In particular,

$$(6.5) \quad \int_{\mathbb{R}^d} f(x)U_+ f(x) dx \leq \int_{\mathbb{R}^d} f(x)Uf(x) dx.$$

Thanks to Lemma 3.1, for all nonnegative functions $f \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ with $\widehat{f} \in L^1(\mathbb{R}^d)$, (6.5) is equivalent to

$$\int_{\mathbb{R}^d} |\widehat{f}(\xi)|^2 \operatorname{Re} \left(\prod_{j=1}^N \frac{1}{1 + \Psi_j(\xi)} \right) d\xi \leq 2^N \int_{\mathbb{R}^d} |\widehat{f}(\xi)|^2 \prod_{j=1}^N \operatorname{Re} \left(\frac{1}{1 + \Psi_j(\xi)} \right) d\xi.$$

Now, given any finite measure μ on the Borel subsets of \mathbb{R}^d , we can replace, in the preceding display, f by $f_\varepsilon \star \mu$, where f_ε is the Gaussian mollifier of (4.6), and obtain

$$\begin{aligned} & \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^2 e^{-(\varepsilon/2)\|\xi\|^2} \operatorname{Re} \left(\prod_{j=1}^N \frac{1}{1 + \Psi_j(\xi)} \right) d\xi \\ & \leq 2^N \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^2 e^{-(\varepsilon/2)\|\xi\|^2} \prod_{j=1}^N \operatorname{Re} \left(\frac{1}{1 + \Psi_j(\xi)} \right) d\xi. \end{aligned}$$

Thanks to (1.3), both integrands are nonnegative. Thus, we can let $\varepsilon \downarrow 0$, and appeal to the Lebesgue monotone convergence theorem to deduce that

$$(2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^2 \operatorname{Re} \left(\prod_{j=1}^N \frac{1}{1 + \Psi_j(\xi)} \right) d\xi \leq 2^N \mathcal{E}_\Psi(\mu).$$

Owing to (6.4), the left-hand side equals $(\mu, \mu)_K$, which completes our proof. \square

PROPOSITION 6.2. *Suppose X is an additive Lévy process on \mathbb{R}^d with Lévy exponent Ψ that satisfies condition (1.3) and X has a one-potential density K . Then, for any compact set $F \subset \mathbb{R}^d$, the following are equivalent:*

- (i) *There exists a finite measure μ on F with $(\mu, \mu)_K < +\infty$.*
- (ii) *$\lambda_d(\mathbf{H}_F) > 0$, where*

$$\mathbf{H}_F = \{a \in \mathbb{R}^d : \mathbb{P}[X(\mathbb{R}_+^N) \cap (\{a\} \oplus F) \neq \emptyset] > 0\}.$$

- (iii) *$\mathbf{H}_F \neq \emptyset$.*

If, in addition, K is almost everywhere positive, then the above is also equivalent to the following:

- (iv) *$\mathbf{H}_F = \mathbb{R}^d$.*

REMARK 6.3. In the classical setting where $N = 1$, this theorem is well known. For instance, when F is a singleton, this proposition was considered first by Orey (1967) and later on by Kesten (1969) and Bretagnolle (1971). The same remark applies to the equivalence of (iv) and (ii). When $F \subset \mathbb{R}^d$ is a general closed set, this result can be found in Bertoin [(1996), Chapter II]. Furthermore, the equivalence of (ii) and (iii) appears in Hawkes [(1979), Theorem 2.1].

PROOF OF PROPOSITION 6.2. We observe that, by (5.9) and by Fubini's theorem,

$$(6.6) \quad \mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N) \ominus F)\} = \int_{\mathbb{R}^d} \mathbb{P}\{X(\mathbb{R}_+^N) \cap [\{a\} \oplus F] \neq \emptyset\} da.$$

Hence, (i) \Leftrightarrow (ii) follows from Theorem 1.5 and Lemma 6.1. It is also clear that (ii) \Rightarrow (iii). To prove (iii) \Rightarrow (ii), note that, for all $s \in \mathbb{R}_+^N$ and all $a \in \mathbb{R}^d$,

$$\begin{aligned} & \mathbb{P}\{X((s, \infty)) \cap [\{a\} \oplus F] \neq \emptyset\} \\ &= \int_{\mathbb{R}^d} \mathbb{P}\{X(\mathbb{R}_+^N) \cap [\{a - b\} \oplus F] \neq \emptyset\} P\{X(s) \in db\}, \end{aligned}$$

where $(s, \infty) = \{t \in \mathbb{R}_+^N; t \succcurlyeq s\}$. We multiply the above by $\exp(-\sum_j s_j)$ and integrate $[ds]$ to obtain

$$(6.7) \quad \begin{aligned} & \int_{\mathbb{R}_+^N} \mathbb{P}\{X((s, \infty)) \cap [\{a\} \oplus F] \neq \emptyset\} \exp\left(-\sum_{j=1}^N s_j\right) ds \\ &= \int_{\mathbb{R}^d} \mathbb{P}\{X(\mathbb{R}_+^N) \cap [\{a - b\} \oplus F] \neq \emptyset\} K(b) db. \end{aligned}$$

If (iii) holds, then for some $a \in \mathbb{R}^d$ the left-hand side of (6.7) is positive. Therefore,

$$\mathbb{P}\{X(\mathbb{R}_+^N) \cap [\{a - b\} \oplus F] \neq \emptyset\} > 0,$$

for b in a set of positive Lebesgue measure, which implies $\lambda_d(H_F) > 0$. This proves the first half of the proposition. Since it is clear that (iv) \Rightarrow (i), it remains to prove (i) \Rightarrow (iv).

If (iv) did not hold, there would exist an $a \in \mathbb{R}^d$ such that the left-hand side of (6.7) would equal to 0. This would then imply that

$$\int_{\mathbb{R}^d} \mathbb{P}\{X(\mathbb{R}_+^N) \cap [\{a - b\} \oplus F] \neq \emptyset\} K(b) db = 0.$$

Since $K > 0$ almost everywhere, we would have

$$\mathbb{P}\{X(\mathbb{R}_+^N) \cap [\{a - b\} \oplus F] \neq \emptyset\} = 0, \quad \lambda_d\text{-almost every } b \in \mathbb{R}^d.$$

Equation (6.6) would then imply that $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N) \ominus F)\} = 0$. Using Theorem 1.5 and Lemma 6.1 again, we would derive a contradiction to (i). We have shown that (i) \Rightarrow (iv), which completes our proof. \square

REMARK 6.4. The almost everywhere positivity of the function K is indispensable, as can be seen by considering a nonnegative stable subordinator X and by letting $F = [-2, -1]$. In this case (iii) clearly does not hold.

In the following, we prove a zero-infinity law for $\lambda_d(X(\mathbb{R}_+^N))$.

PROPOSITION 6.5. *Suppose X is an additive Lévy process on \mathbb{R}^d with Lévy exponent Ψ that satisfies condition (1.3) and has an a.e. positive 1-potential density K . Then*

$$\lambda_d(X(\mathbb{R}_+^N)) \in \{0, +\infty\}, \quad \mathbb{P}\text{-a.s.}$$

PROOF. Assuming that $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} < \infty$, we first show that the value of this expectation is, in fact, zero. Bearing this goal in mind, we note that, for any $n > 0$,

$$\begin{aligned} \mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} &\geq \mathbb{E}\{\lambda_d(X([0, n]^N))\} + \mathbb{E}\{\lambda_d(X((n, \infty)^N))\} \\ &\quad - \mathbb{E}\{\lambda_d[X([0, n]^N) \cap X((n, \infty)^N)]\} \\ &= \mathbb{E}\{\lambda_d(X([0, n]^N))\} + \mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} \\ &\quad - \mathbb{E}\{\lambda_d[X([0, n]^N) \cap X'(\mathbb{R}_+^N)]\}, \end{aligned}$$

where X' is an independent copy of X . Consequently, we see that if $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} < \infty$,

$$\mathbb{E}\{\lambda_d(X([0, n]^N))\} \leq \mathbb{E}\{\lambda_d[X([0, n]^N) \cap X'(\mathbb{R}_+^N)]\}.$$

Let $n \uparrow \infty$ to see that, as long as $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} < \infty$,

$$\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} \leq \mathbb{E}\{\lambda_d[X(\mathbb{R}_+^N) \cap X'(\mathbb{R}_+^N)]\}.$$

Define $\varphi(a) = \mathbb{P}\{a \in X(\mathbb{R}_+^N)\}$, and note that the above is equivalent to

$$\int_{\mathbb{R}^d} \varphi(a) da \leq \int_{\mathbb{R}^d} \varphi^2(a) da.$$

Since $0 \leq \varphi(a)(1 - \varphi(a)) \leq 1$ for all $a \in \mathbb{R}^d$, we have

$$\varphi(a) \in \{0, 1\}, \quad \lambda_d\text{-almost every } a \in \mathbb{R}^d.$$

Consequently, if $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\}$ is finite,

$$(6.8) \quad \mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} = \lambda_d(\varphi^{-1}\{1\}).$$

It follows from Proposition 6.2 that either $\varphi(a) = 0$ for all $a \in \mathbb{R}^d$ or $\varphi(a) > 0$ for all $a \in \mathbb{R}^d$. This means that $\lambda_d(\varphi^{-1}\{1\}) = 0$; for, otherwise, $\varphi^{-1}\{1\} = \mathbb{R}^d$, which

has infinite λ_d -measure, and this would contradict (6.8). In other words, we have demonstrated that

$$\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} < \infty \implies \lambda_d(\varphi^{-1}\{1\}) = 0 \implies \mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} = 0.$$

We now “remove the expectation” from this statement and finish our proof. Suppose that $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} > 0$ (which means that $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} = \infty$), and note that, for any $\nu > 0$,

$$\lambda_d(X(\mathbb{R}_+^N)) \geq \sup_{n \geq 0} \lambda_d(X([n, n + \nu]^N)) = \sup_{n \geq 0} \Gamma_n^\nu.$$

Since $\Gamma_0^\nu, \Gamma_{1+\nu}^\nu, \Gamma_{1+2\nu}^\nu, \dots$ are i.i.d., by the Borel–Cantelli lemma, for any $\nu > 0$,

$$\lambda_d(X(\mathbb{R}_+^N)) \geq \mathbb{E}\{\Gamma_0^\nu\} = \mathbb{E}\{\lambda_d(X([0, \nu]^N))\}, \quad \mathbb{P}\text{-a.s.}$$

We can let $\nu \uparrow \infty$ along a sequence of rational numbers to deduce from $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} = \infty$ that $\lambda_d(X(\mathbb{R}_+^N)) = \infty$, a.s. \square

7. Bessel–Riesz capacities. The α -dimensional Bessel–Riesz (sometimes only Riesz) energies and capacities on \mathbb{R}^d are those that correspond to $K = \mathcal{R}_{(\alpha)}$, where

$$(7.1) \quad \mathcal{R}_{(\alpha)}(a) = \begin{cases} 1, & \text{if } \alpha < 0, \\ \ln(1/\|a\|), & \text{if } \alpha = 0, \\ \|a\|^{-\alpha}, & \text{if } \alpha > 0, \end{cases} \quad \forall a \in \mathbb{R}^d.$$

In this case we will write $(\mu, \mu)_{(\alpha)}$, in place of the more cumbersome $(\mu, \mu)_{\mathcal{R}_{(\alpha)}}$, and write $C_{(\alpha)}$ for the corresponding capacity.

There are deep connections between α -dimensional Bessel–Riesz capacities and ordinary one-parameter Lévy processes when $0 < \alpha \leq 2$. In this section, we show that, by considering additive Lévy processes, one can have a probabilistic interpretation of sets of positive α -dimensional capacity for any $\alpha > 0$.

EXAMPLE 7.1. Suppose $B = B_1 \oplus \dots \oplus B_N$ is additive Brownian motion in \mathbb{R}^d . That is, B_i ’s are independent d -dimensional Brownian motions. Then, by (6.2), it is easy to see that the 1-potential density of B is

$$K(a) = \frac{1}{(2\pi)^{d/2}(N-1)!} \int_0^\infty e^{-t-\|a\|^2/(2t)} t^{-d/2+N-1} dt \quad \forall a \in \mathbb{R}^d.$$

This calculation only requires the elementary fact that

$$\lambda_k(\{q \in \mathbb{R}_+^k : q_1 + \dots + q_k \leq x\}) = \frac{1}{k!} x^k \quad \forall x \geq 0, k = 1, 2, \dots,$$

which, itself, follows from symmetry considerations. Furthermore, it is a simple matter to check that

$$\lim_{a \rightarrow 0} \frac{K(a)}{\mathcal{R}_{(d-2N)}(a)} = \frac{\Lambda(d/2 - N)}{(2\pi)^{d/2}(N - 1)!},$$

where $\mathcal{R}_{(\alpha)}$ is defined in the Riesz kernel of (7.1), and for all $x \in \mathbb{R}$,

$$\Lambda(x) = 2^{x+} \Gamma(|x|),$$

where $x_+ = \max(x, 0)$. Consequently, we can deduce that, for any compact set $F \subset \mathbb{R}^d$, there are two constants A_1 and A_2 such that, for all $x \in F \oplus F$,

$$A_1 \mathcal{R}_{(d-2N)}(x) \leq K(x) \leq A_2 \mathcal{R}_{(d-2N)}(x).$$

This, Lemma 6.1 and Theorem 1.5, together, combine to show that $B(\mathbb{R}_+^N) \oplus F$ can have positive Lebesgue measure if and only if $C_{(d-2N)}(F) > 0$.

With a little more work, and motivated by this example, we can find additive Lévy processes that correspond to *any* Bessel–Riesz capacity of interest. Recall that $X_1 \oplus \dots \oplus X_N$ is *additive stable* of index $\alpha \in (0, 2]$, if X_1, \dots, X_N are independent isotropic stable processes with index α each.

THEOREM 7.2. *Suppose $X = X_1 \oplus \dots \oplus X_N$ is an additive stable process of index $\alpha \in (0, 2]$, and in \mathbb{R}^d . Then, for any compact set $F \subset \mathbb{R}^d$,*

$$\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N) \oplus F)\} > 0 \iff C_{(d-\alpha N)}(F) > 0.$$

REMARK 7.3. Upon varying $d, N \in \mathbb{N}$ and $\alpha \in (0, 2]$, we see that this theorem associates an additive Lévy process to any Bessel–Riesz capacity, including those with dimension greater than 2.

Theorem 7.2 follows from Theorem 1.5 and the arguments of Example 7.1, once we establish the following.

PROPOSITION 7.4. *Let X denote an N -parameter additive stable process in \mathbb{R}^d . Then X has a one-potential density K whose asymptotics at the origin are described by the following:*

$$(7.2) \quad \lim_{a \rightarrow 0} \frac{K(a)}{\mathcal{R}_{(d-N\alpha)}(a)} = C(\alpha, d, N),$$

where $\mathcal{R}_{(\cdot)}$ is the Riesz kernel of (7.1), and $C(\alpha, d, N)$ is a positive and finite constant depending on α, d and N only. Moreover, there exists a positive constant $\tilde{C} = \tilde{C}(\alpha, d, N)$ such that

$$(7.3) \quad K(a) \leq \tilde{C} \mathcal{R}_{(d-N\alpha)}(a) \quad \forall a \in \mathbb{R}^d.$$

PROOF. In light of Example 7.1, we can assume, without loss generality, that $0 < \alpha < 2$.

We denote the density function of $X_1(1)$ by $p_1(x)$. Here, p_1 is scaled as

$$e^{-\|\xi\|^\alpha} = \int_{\mathbb{R}^d} e^{ix \cdot \xi} p_1(x) dx.$$

It is possible to show that $p_1(x)$ is a continuous and strictly positive function on \mathbb{R}^d that is isotropic; that is, it depends on x only through $\|x\|$.

Direct calculations reveal that the 1-potential density of X is

$$(7.4) \quad K(a) = \frac{1}{(N-1)!} \int_0^\infty t^{N-1-d/\alpha} e^{-t} p_1(at^{-1/\alpha}) dt \quad \forall a \in \mathbb{R}^d.$$

On the other hand, by using Bochner’s subordination, we can write

$$(7.5) \quad p_1(x) = \|x\|^{-d} \int_0^\infty v\left(\frac{s}{\|x\|^2}\right) g^{(\alpha/2)}(s) ds,$$

where the function v is defined as

$$v(s) = (4\pi s)^{-d/2} \exp\left(-\frac{1}{4s}\right),$$

$g^{(\alpha/2)}(s)$ is the density function of the random variable $\tau(1)$, and where $\tau = \{\tau(t), t \geq 0\}$ is a stable subordinator of index $\frac{1}{2}\alpha$ [cf., e.g., Bendikov (1994)]. It should be recognized that $a \mapsto K(a)$ is isotropic and strictly decreasing in $\|a\|$.

It follows from (7.4) and (7.5), combined with Fubini’s theorem, that

$$(7.6) \quad \begin{aligned} K(a) &= \frac{\|a\|^{-d}}{(N-1)!} \int_0^\infty \int_0^\infty t^{N-1} e^{-t} v\left(\frac{st^{2/\alpha}}{\|a\|^2}\right) g^{(\alpha/2)}(s) ds dt \\ &= \frac{\alpha/2}{(4\pi)^{d/2}(N-1)!} \|a\|^{\alpha N-d} \int_0^\infty s^{-\alpha N/2} g^{(\alpha/2)}(s) ds \\ &\quad \times \int_0^\infty u^{(\alpha N-d)/2-1} \exp\left(-\frac{\|a\|^\alpha}{s^{\alpha/2}} u^{\alpha/2}\right) \exp\left(-\frac{1}{4u}\right) du. \end{aligned}$$

When $d - \alpha N > 0$, (7.6) and the monotone convergence theorem imply

$$\begin{aligned} &\lim_{a \rightarrow 0} \frac{K(a)}{\mathcal{R}_{(d-\alpha N)}(a)} \\ &= \frac{\alpha/2}{(4\pi)^{d/2}(N-1)!} \int_0^\infty s^{-\alpha N/2} g^{(\alpha/2)}(s) ds \\ &\quad \times \int_0^\infty u^{(\alpha N-d)/2-1} \exp\left(-\frac{1}{4u}\right) du. \end{aligned}$$

Hence, when $d - \alpha N > 0$, we can identify the constant $C(\alpha, d, N)$ in (7.2) as

$$C(\alpha, d, N) = \frac{\alpha 2^{d-\alpha N-1}}{(4\pi)^{d/2}(N-1)!} \Gamma\left(\frac{1}{2}(d-\alpha N)\right) \int_0^\infty s^{-\alpha N/2} g^{(\alpha/2)}(s) ds.$$

In the above, note that

$$\begin{aligned}
 (7.7) \quad \int_0^\infty s^{-\alpha N/2} g^{(\alpha/2)}(s) ds &= \mathbb{E}[\tau(1)^{-\alpha N/2}] \\
 &= \int_0^\infty \mathbb{P}\{\tau(1) \leq x^{-2/(\alpha N)}\} dx < +\infty,
 \end{aligned}$$

by a well-known estimate for $\mathbb{P}\{\tau(1) \leq \varepsilon\}$ as $\varepsilon \rightarrow 0$ [see, e.g., Hawkes (1971) or Bertoin (1996), page 88].

If $d - \alpha N = 0$, we split the last integral in (7.6) as

$$(7.8) \quad \int_0^{\|a\|^{-1}} (\dots) du + \int_{\|a\|^{-1}}^\infty (\dots) du = I_1(a) + I_2(a).$$

We note that

$$\begin{aligned}
 &\exp\left(-\frac{\|a\|^{\alpha/2}}{s^{\alpha/2}}\right) \int_0^{\|a\|^{-1}} u^{-1} \exp\left(-\frac{1}{4u}\right) du \\
 &\leq I_1(a) \leq \int_0^{\|a\|^{-1}} u^{-1} \exp\left(-\frac{1}{4u}\right) du,
 \end{aligned}$$

and, thanks to l'Hôpital's rule,

$$\lim_{a \rightarrow 0} \frac{\int_0^{\|a\|^{-1}} u^{-1} \exp(-1/(4u)) du}{\log \|a\|^{-1}} = 1.$$

Thus,

$$(7.9) \quad \lim_{a \rightarrow 0} \frac{I_1(a)}{\log \|a\|^{-1}} = 1.$$

On the other hand,

$$\begin{aligned}
 &\exp\left(-\frac{\|a\|}{4}\right) \int_{\|a\|^{-1}}^\infty u^{-1} \exp\left(-\frac{\|a\|^\alpha}{s^{\alpha/2}} u^{\alpha/2}\right) du \\
 &\leq I_2(a) \leq \int_{\|a\|^{-1}}^\infty u^{-1} \exp\left(-\frac{\|a\|^\alpha}{s^{\alpha/2}} u^{\alpha/2}\right) du,
 \end{aligned}$$

and, by using l'Hôpital's rule again, we obtain

$$\begin{aligned}
 (7.10) \quad \lim_{a \rightarrow 0} \frac{I_2(a)}{\log \|a\|^{-1}} &= \lim_{a \rightarrow 0} \frac{1}{\log \|a\|^{-1}} \int_{\|a\|^{-1}}^\infty u^{-1} \exp\left(-\frac{\|a\|^\alpha}{s^{\alpha/2}} u^{\alpha/2}\right) du \\
 &= \lim_{a \rightarrow 0} \left[-\exp\left(-\frac{\|a\|^{\alpha/2}}{s^{\alpha/2}}\right) \right. \\
 &\quad \left. + \frac{\alpha \|a\|^\alpha}{s^{\alpha/2}} \int_{\|a\|^{-1}}^\infty u^{-1+\alpha/2} \exp\left(-\frac{\|a\|^\alpha}{s^{\alpha/2}} u^{\alpha/2}\right) du \right] \\
 &= 1.
 \end{aligned}$$

It follows from (7.6) and (7.8)–(7.10), combined with the Lebesgue dominated convergence theorem [the above upper bounds for $I_1(a)$ and $I_2(a)$ are used here], that when $d - \alpha N = 0$,

$$\lim_{a \rightarrow 0} \frac{K(a)}{\mathcal{R}_{(0)}(a)} = \frac{\alpha}{(4\pi)^{d/2}(N-1)!} \int_0^\infty s^{-\alpha N/2} g^{(\alpha/2)}(s) ds.$$

In case $d - \alpha N < 0$, (7.6) tells us that we only need to show that the following limit exists:

$$\lim_{a \rightarrow 0} \|a\|^{\alpha N - d} \int_0^\infty u^{(\alpha N - d)/2 - 1} \exp\left(-\frac{\|a\|^\alpha}{s^{\alpha/2}} u^{\alpha/2}\right) \exp\left(-\frac{1}{4u}\right) du.$$

After changing variables and appealing to the monotone convergence theorem, we can see that the above limit equals

$$\int_0^\infty v^{(\alpha N - d)/2 - 1} \exp\left(-\frac{v^{\alpha/2}}{s^{\alpha/2}}\right) dv.$$

Hence, in this case, we apply (7.6), and change variables once more, to show that (7.2) holds with

$$C(\alpha, d, N) = \frac{\Gamma(N - d/\alpha)}{(4\pi)^{d/2}(N-1)!} \int_0^\infty s^{-d/2} g^{(\alpha/2)}(s) ds,$$

where, as in (7.7), the last integral is finite.

Finally, the inequality (7.3) follows readily by adapting the aforementioned arguments. For example, when $d - \alpha N > 0$, it follows from (7.6) that

$$\begin{aligned} \frac{K(a)}{\mathcal{R}_{(d - N\alpha)}(a)} &\leq \frac{\alpha/2}{(4\pi)^{d/2}(N-1)!} \int_0^\infty s^{-\alpha N/2} g^{(\alpha/2)}(s) ds \\ &\quad \times \int_0^\infty u^{(\alpha N - d)/2 - 1} \exp\left(-\frac{1}{4u}\right) du. \end{aligned}$$

We omit the other two cases and declare the proof of Proposition 7.4 complete. □

8. Proof of Theorem 1.6. We will use Frostman’s theorem of potential theory [cf. Kahane (1985), Chapter 10, or Carleson (1983)]. Recall that the latter states that, for any Borel set $G \subset \mathbb{R}^d$, the capacity and the Hausdorff dimensions of G agree. That is,

$$(8.1) \quad \dim(G) = \sup\{\gamma > 0 : \mathbf{C}_{(\gamma)}(G) > 0\},$$

where $\mathbf{C}_{(\gamma)}$ is the Bessel–Riesz capacity of Section 7, and $\sup \emptyset = 0$.

Now we introduce an M -parameter additive stable process Y in \mathbb{R}^d whose index is $\alpha \in (0, 2]$. The process Y is totally independent of X , and we will determine the

constants M and α shortly. Note that $X \oplus Y$ is an $(N + M)$ -parameter additive Lévy process in \mathbb{R}^d whose Lévy exponent $\Phi = (\Phi_1, \dots, \Phi_{N+M})$ is given by

$$\Phi_j(\xi) = \begin{cases} \Psi_j(\xi), & \text{if } j = 1, \dots, N, \\ \frac{1}{2}\|\xi\|^\alpha, & \text{if } j = N + 1, \dots, N + M. \end{cases}$$

Clearly,

$$\int_{\mathbb{R}^d} \prod_{j=1}^{N+M} \operatorname{Re}\left(\frac{1}{1 + \Phi_j(\xi)}\right) d\xi < +\infty \iff \mathcal{J}^{(M\alpha)} < +\infty,$$

where, for all $\gamma \in \mathbb{R}$,

$$\mathcal{J}^{(\gamma)} = \int_{\xi \in \mathbb{R}^d : \|\xi\| > 1} \prod_{j=1}^N \operatorname{Re}\left(\frac{1}{1 + \Psi_j(\xi)}\right) \|\xi\|^{-\gamma} d\xi.$$

Thus, we can apply Theorem 1.1 to the process $X \oplus Y$ and see that

$$\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N) \oplus Y(\mathbb{R}_+^M))\} > 0 \iff \mathcal{J}^{(M\alpha)} < +\infty.$$

On the other hand, we can also apply Theorem 7.2, conditionally on $F = X(\mathbb{R}_+^N)$, to deduce that

$$\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N) \oplus Y(\mathbb{R}_+^M))\} > 0 \iff \mathbb{E}\{\mathbf{C}_{(d-M\alpha)}(X(\mathbb{R}_+^N))\} > 0.$$

We combine the latter two displays to obtain

$$\mathbb{E}\{\mathbf{C}_{(d-M\alpha)}(X(\mathbb{R}_+^N))\} > 0 \iff \mathcal{J}^{(M\alpha)} < +\infty.$$

Consequently, when $\mathcal{J}^{(M\alpha)} = +\infty$, $\mathbf{C}_{(d-M\alpha)}(X(\mathbb{R}_+^N)) = 0$, \mathbb{P} -almost surely. From Frostman’s theorem [(8.1)], we deduce that $\mathcal{J}^{(M\alpha)} = +\infty$ implies that, \mathbb{P} -a.s., $\dim(X(\mathbb{R}_+^N)) \leq d - M\alpha$. On the other hand, we can choose $M \in \{1, 2, \dots\}$ and rational $\alpha_1, \alpha_2, \dots \in (0, 2]$ such that $M\alpha_j \uparrow \eta$; this shows that $\dim(X(\mathbb{R}_+^N)) \leq d - \eta$, \mathbb{P} -almost surely. In particular, if $\eta = d$, then $\dim(X(\mathbb{R}_+^N)) = 0$, \mathbb{P} -almost surely, and this constitutes half of our theorem. For the other half, we use the same argument, but quantitatively.

For the converse half, we only need to consider the case when $\eta < d$. With this in mind, choose $\alpha \in (0, 2]$ and $M \in \{1, 2, \dots\}$ such that $\eta < M\alpha < d$. Thus, we can deduce from the preceding paragraph that

$$\mathcal{J}^{(M\alpha)} < +\infty.$$

We now recall the killed occupation measure \mathbb{O}_κ from (2.1) and (5.1). This is a Borel probability measure carried by $X(\mathbb{R}_+^N)$, and we claim that, as long as $0 < d - \alpha M < d - \eta$,

$$(8.2) \quad (\mathbb{O}_\kappa, \mathbb{O}_\kappa)_{(d-M\alpha)} < +\infty, \quad \mathbb{P}\text{-a.s.},$$

where $(\mathbb{O}_\kappa, \mathbb{O}_\kappa)_{(\gamma)}$ is the γ -dimensional Bessel–Riesz energy of \mathbb{O}_κ as defined in Section 7. Together with Frostman’s theorem [(8.1)], this shows that with probability 1, $\dim(X(\mathbb{R}_+^N)) \geq d - M\alpha$. This is the key part of our proof, since we can approximate η from above arbitrarily well by numbers of the form $M\alpha$ ($M \in \{1, 2, \dots\}$, $\alpha \in (0, 2] \cap \mathbb{Q}$). In this way, we deduce that, with probability 1, $\dim(X(\mathbb{R}_+^N)) \geq d - \eta$, as asserted.

At this point, we only need to establish (8.2). For this purpose, recall the process Y , as above, and consider its Lévy exponent $\Lambda = (\Lambda_1, \dots, \Lambda_M)$, where $\Lambda_i(\xi) = \frac{1}{2}\|\xi\|^\alpha$, $i = 1, \dots, M$. The process Y has a 1-potential density K whose asymptotics are described by Proposition 7.4. After applying Lemma 6.1 to Y , we deduce that $(\mathbb{O}_\kappa, \mathbb{O}_\kappa)_K = \mathcal{E}_\Lambda(\mathbb{O}_\kappa)$, \mathbb{P} -almost surely. In particular, Lemma 5.2 gives

$$\begin{aligned} \mathbb{E}\{(\mathbb{O}_\kappa, \mathbb{O}_\kappa)_K\} &= (2\pi)^{-d} \int_{\mathbb{R}^d} |\widehat{\mathbb{O}_\kappa}(\xi)|^2 \cdot \left\{1 + \frac{1}{2}\|\xi\|^\alpha\right\}^{-M} d\xi \\ &= (2\pi)^{-d} \int_{\mathbb{R}^d} \prod_{j=1}^N \operatorname{Re}\left(\frac{1}{1 + \Psi_j(\xi)}\right) \cdot \left\{1 + \frac{1}{2}\|\xi\|^\alpha\right\}^{-M} d\xi, \end{aligned}$$

which is finite, since $\mathcal{J}^{(M\alpha)} < +\infty$. Therefore, we have found a random measure \mathbb{O}_κ on the random set $X(\mathbb{R}_+^N)$ such that, with probability 1, $(\mathbb{O}_\kappa, \mathbb{O}_\kappa)_K < +\infty$. Thanks to Proposition 7.4, there is a positive and finite constant C' , depending on α , d and N only, such that

$$\mathcal{R}_{(d-N\alpha)}(a) \leq C'K(a) \quad \forall a \in \mathbb{R}^d \text{ with } \|a\| \leq 1.$$

Hence

$$(\mathbb{O}_\kappa, \mathbb{O}_\kappa)_{(d-M\alpha)} \leq 1 + C' \times (\mathbb{O}_\kappa, \mathbb{O}_\kappa)_K < +\infty.$$

This verifies (8.2), whence Theorem 1.6 follows.

9. Concluding remarks. A number of interesting questions remain unresolved, some of which are listed below.

QUESTION 9.1. Do Theorems 1.1 and 1.5 hold for all additive Lévy processes? One only needs to worry about the necessity since in both theorems the sufficiency has already been shown to hold generally.

A possible approach for proving Theorem 1.1 without condition (1.3) is as follows: In light of Lemma 5.2 and the upper bound in Theorem 2.1, it suffices to show that

$$(9.1) \quad \exists \mu \in \mathcal{P}(E) : \|\mu\|_e < +\infty \implies \|\kappa\|_e < +\infty.$$

When $\|\cdot\|_e$ is an energy norm based on a positive definite convolution kernel, one can prove such a result by appealing to simple Fourier analytical arguments. In the present general setting, however, we do not know how to proceed. In the Appendix,

we derive an analytical estimate that may be of independent interest and that barely falls short of settling this open problem by way of verifying the preceding display.

QUESTION 9.2. A simpler, but still interesting, open problem is to find a necessary and sufficient condition for $X(\mathbb{R}_+^N) \oplus F$ to have positive Lebesgue measure with positive probability when X_i 's are independent (but, otherwise, arbitrary) subordinators. Equivalently, we ask for a necessary and sufficient condition for the existence of local times of N -parameter additive subordinators without a condition such as (1.3).

QUESTION 9.3. In light of Theorem 1.6, it would be interesting to determine an exact Hausdorff measure function that gauges the size of $X([0, 1]^N)$. For stable sheets and two-parameter additive subordinators, related results can be found in Ehm (1981) and Hu (1994).

APPENDIX

In this Appendix we present a possible alternative approach for proving Theorem 1.1 without condition (1.3) that involves an estimate that may be of independent analytical interest. As we mentioned in Section 9, the key is to prove (9.1). In the present general setting, we are only able to verify a partial derivation (cf. Proposition A.3 and Remark A.4).

Given any measure μ on \mathbb{R}_+^N , and given $s \in \mathbb{R}^N$, write

$$\mu_{(s)}(\cdot) = \mu(\cdot + s)$$

for the s -shift of μ . We note that $\mu_{(s)}$ need not be a measure on \mathbb{R}_+^N although μ is assumed to be.

LEMMA A.1 (Shift invariance of energy norm). *The map $\mu \mapsto \|\mu\|_e$ is shift invariant in the sense that, whenever μ and $\mu_{(s)}$ are both finite measures on \mathbb{R}_+^N ,*

$$\|\mu\|_e = \|\mu_{(s)}\|_e.$$

PROOF. This follows immediately from our computation of $\|\mu\|_e$ in Lemma 2.4. \square

Next, we prove that convolutions reduce the norm. To be more precise, we have the following.

LEMMA A.2 (Norm reduction of convolutions). *Suppose φ is a probability density function on \mathbb{R}^N such that $\varphi \star \mu$ and μ are both in $\mathcal{P}(\mathbb{R}_+^N)$. Then*

$$\|\varphi \star \mu\|_e \leq \|\mu\|_e.$$

PROOF. We write $\varphi \star \mu$ for both the measure and its density with respect to Lebesgue measure λ_N . In this way we can write the corresponding occupation measure as

$$\mathbb{O}_{\varphi \star \mu}(f) = \int_{\mathbb{R}_+^N} f(X(s)) \varphi \star \mu(s) ds.$$

Using Fubini’s theorem, twice in succession, we obtain

$$\begin{aligned} \widehat{\mathbb{O}_{\varphi \star \mu}}(\xi) &= \int_{\mathbb{R}_+^N} e^{i\xi \cdot X(s)} ds \int_{\mathbb{R}_+^N} \varphi(s-t) \mu(dt) \\ &= \int_{\mathbb{R}_+^N} \mu(dt) \int_{\mathbb{R}_+^N \ominus \{t\}} e^{i\xi \cdot X(v+t)} \varphi(v) dv \\ &= \int_{\mathbb{R}^N} \varphi(v) dv \int_{\mathbb{R}_+^N \ominus \{v\}} e^{i\xi \cdot X(v+t)} \mu(dt) \\ &= \int_{\mathbb{R}^N} \widehat{\mathbb{O}_{\mu(v)}}(\xi) \varphi(v) dv. \end{aligned}$$

To this, we apply the Cauchy–Schwarz inequality twice as follows:

$$\begin{aligned} \|\varphi \star \mu\|_e^2 &= (2\pi)^{-d} \int_{\mathbb{R}^d} \iint_{\mathbb{R}^N \times \mathbb{R}^N} \mathbb{E}\left\{ \widehat{\mathbb{O}_{\mu(u)}}(\xi) \overline{\widehat{\mathbb{O}_{\mu(v)}}(\xi)} \right\} \varphi(u) \varphi(v) du dv d\xi \\ &\leq (2\pi)^{-d} \int_{\mathbb{R}^d} \left| \int_{\mathbb{R}^N} \sqrt{\mathbb{E}\{|\widehat{\mathbb{O}_{\mu(v)}}(\xi)|^2\}} \varphi(v) dv \right|^2 d\xi \\ &\leq (2\pi)^{-d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^N} \mathbb{E}\{|\widehat{\mathbb{O}_{\mu(v)}}(\xi)|^2\} \varphi(v) dv d\xi \\ &= \|\mu\|_e^2, \end{aligned}$$

thanks to Lemma A.1. \square

We conclude this Appendix by showing that if there are any probability measures of finite norm on a given compact set F , Lebesgue measure on F is also of finite norm.

PROPOSITION A.3. *Suppose $F \subset \mathbb{R}_+^N$ is compact. Then*

$$\|\lambda_N \lfloor_F\|_e \leq 2^N \sup_{t \in F} |t|^N \inf_{\mu \in \mathcal{P}(F)} \|\mu\|_e,$$

where $\lambda_N \lfloor_F$ denotes the restriction of λ_N to F .

PROOF. First, we suppose that F is a compact subset of $(0, \infty)^N$. At the end of our proof, we show how this condition can be removed.

Since $F \subset (0, \infty)^N$, its closed ε -enlargement (written as F^ε) is a subset of \mathbb{R}_+^N for all $\varepsilon > 0$ small enough.

For all $\varepsilon > 0$, define the function φ_ε , on \mathbb{R}^N , by

$$\varphi_\varepsilon(r) = (2\varepsilon)^{-N} \mathbb{1}_{\mathcal{B}(0,\varepsilon)}(r) \quad \forall r \in \mathbb{R}^N,$$

where

$$\mathcal{B}(s, r) = \{t \in \mathbb{R}^N : |t - s| \leq r\} \quad \forall s \in \mathbb{R}^N, r > 0,$$

and $|t| = \max_{1 \leq i \leq N} |t_i|$ is the ℓ^∞ -norm on \mathbb{R}^N . We note the following logical sequence:

$$\begin{aligned} \mu \in \mathcal{P}(F) &\implies \varphi_\varepsilon \star \mu \in \mathcal{P}(F^\varepsilon) \\ &\implies \forall \varepsilon > 0 \text{ small, } \varphi_\varepsilon \star \mu \in \mathcal{P}(\mathbb{R}_+^N) \\ &\implies \forall \varepsilon > 0 \text{ small, } \|\varphi_\varepsilon \star \mu\|_e \leq \|\mu\|_e, \end{aligned}$$

the last inequality following from the norm reduction of convolutions (cf. Lemma A.2). For each $\eta > 0$, let f_η denote the density of $\eta^{1/2}$ times a d -dimensional vector of independent standard Gaussians [cf. (4.6)]. We note that, for all $\varepsilon > 0$ sufficiently small,

$$\mathbb{O}_{\varphi_\varepsilon \star \mu}(f_\eta) = (2\varepsilon)^{-N} \int f_\eta(X(s)) \mu(\mathcal{B}(s, \varepsilon)) ds.$$

[The only reason for our insisting on the smallness of ε is to ensure that $\varphi_\varepsilon \star \mu \in \mathcal{P}(\mathbb{R}_+^N)$. Of course, here, “small” means “small enough to ensure that $F^\varepsilon \subset \mathbb{R}_+^N$.”] We will use this formula, and a covering argument, to obtain a simple bound.

For any compact set $K \subset \mathbb{R}_+^N$, and for all $\varepsilon > 0$, let $\mathcal{N}_K(\varepsilon)$ denote the minimum number of ℓ^∞ -balls of radius ε needed to cover K . \mathcal{N}_K is sometimes called the *metric entropy* of K . Plainly, if $\varepsilon > 0$ is fixed but sufficiently small, we can find $s^1, \dots, s^{\mathcal{N}_F(\varepsilon)} \in \mathbb{R}_+^N$ such that $\mathcal{B}(s^j, \varepsilon) \subset \mathbb{R}_+^N$ and $\bigcup_{j=1}^{\mathcal{N}_F(\varepsilon)} \mathcal{B}(s^j, \varepsilon) \supseteq F$. Consequently, $\sum_{j=1}^{\mathcal{N}_F(\varepsilon)} \mu(\mathcal{B}(s^j, \varepsilon)) \geq \mu(F) = 1$. We can deduce the existence of a point $s^* \in \mathbb{R}_+^N$ such that $\mu(\mathcal{B}(s^*, \varepsilon)) \geq [\mathcal{N}_F(\varepsilon)]^{-1}$. (Warning: s^* may depend on ε .) On the other hand, for all $s \in \mathcal{B}(s^*, \varepsilon)$, $\mathcal{B}(s, 2\varepsilon) \supseteq \mathcal{B}(s^*, \varepsilon)$, thanks to the triangle inequality. In other words, we have shown that for

$$\inf_{s \in \mathcal{B}(s^*, \varepsilon)} \mu(\mathcal{B}(s, 2\varepsilon)) \geq \frac{1}{\mathcal{N}_F(\varepsilon)}.$$

The display preceding the above, then, shows that

$$\begin{aligned} \mathbb{O}_{\varphi_{2\varepsilon} \star \mu}(f_\eta) &\geq \frac{1}{(4\varepsilon)^N \mathcal{N}_F(\varepsilon)} \int_{\mathcal{B}(s^*, \varepsilon)} f_\eta(X(s)) ds \\ &= \frac{1}{(4\varepsilon)^N \mathcal{N}_F(\varepsilon)} \mathbb{O}_{\lambda_N \lfloor_{\mathcal{B}(s^*, \varepsilon)}}(f_\eta). \end{aligned}$$

In other words,

$$\mathbb{E}_{\lambda_d} \{ |\mathbb{O}_{\varphi_{2\varepsilon} \star \mu}(f_\eta)|^2 \} \geq \frac{1}{(4\varepsilon)^{2N} [\mathcal{N}_F(\varepsilon)]^2} \mathbb{E}_{\lambda_d} \{ |\mathbb{O}_{\lambda_N \lfloor_{\mathcal{B}(s^*, \varepsilon)}}(f_\eta)|^2 \}.$$

We apply Lemma 3.5 to see that for any finite measure ν , on \mathbb{R}_+^N ,

$$(A.1) \quad \mathbb{E}\lambda_d\{|\mathbb{O}_\nu(f_\eta)|^2\} \uparrow \|\nu\|_e^2 \quad \text{as } \eta \downarrow 0.$$

Thus, we have shown that, for all $\varepsilon > 0$ sufficiently small,

$$\|\varphi_{2\varepsilon} \star \mu\|_e^2 \geq \frac{1}{(4\varepsilon)^{2N}[\mathcal{N}_F(\varepsilon)]^2} \|\lambda_N \lfloor_{\mathcal{B}(s^*, \varepsilon)}\|_e^2.$$

Since convolutions decrease the energy norm, the first term is bounded above by $\|\mu\|_e^2$ (cf. Lemma A.1). By shift invariance, we then conclude that, for any ℓ^∞ -ball B of radius ε ,

$$\|\mu\|_e^2 \geq \frac{1}{(4\varepsilon)^{2N}[\mathcal{N}_F(\varepsilon)]^2} \|\lambda_N \lfloor_B\|_e^2,$$

as long as $\varepsilon > 0$ is sufficiently small. However, $\|\cdot\|_e$ is a seminorm (Lemma 5.1). Thus

$$(A.2) \quad \begin{aligned} \|\lambda_N \lfloor_F\|_e &\leq \sum_{j=1}^{\mathcal{N}_F(\varepsilon)} \|\lambda_N \lfloor_{\mathcal{B}(s^j, \varepsilon)}\|_e \\ &\leq 4^N \varepsilon^N |\mathcal{N}_F(\varepsilon)|^2 \cdot \|\mu\|_e, \end{aligned}$$

for any $\varepsilon > 0$ that makes $F^\varepsilon \subset \mathbb{R}_+^N$. We now claim that this holds for all $\varepsilon > 0$ and remove the assumption that $F \subset (0, \infty)^N$ in one sweep.

For any real number $c > 0$, consider the set $\langle c \rangle + F = \{\langle c \rangle + s; s \in F\}$, where $\langle c \rangle$ is the N -vector all of whose coordinates are c . By shift invariance, $\lambda_N \lfloor_F$ and $\lambda_N \lfloor_{\langle c \rangle + F}$ have the same energy norm (Lemma A.1). Furthermore, $\mathcal{N}_{\langle c \rangle + F} = \mathcal{N}_F$. Thus, if we only know that $F \subset \mathbb{R}_+^N$, by considering $\langle c \rangle + F$ in place of F in (A.2), we arrive at the following:

$$\|\lambda_N \lfloor_F\|_e \leq 4^N \varepsilon^N |\mathcal{N}_F(\varepsilon)|^2 \cdot \|\mu\|_e,$$

whenever $(\langle c \rangle + F)^\varepsilon \subset \mathbb{R}_+^N$, which holds when $c > \varepsilon$. Since the above inequality is independent of c , we can deduce that (A.2) holds for all compact sets $F \subset \mathbb{R}_+^N$ and for all $\varepsilon > 0$. Let ε equal the ℓ^∞ -radius of F and note that, for this choice of ε , $\mathcal{N}_F(\varepsilon) = 1$, while $\varepsilon^N \leq 2^{-N} \sup_{t \in F} |t|^N$. \square

REMARK A.4. Proposition A.3 comes very close to showing that Theorem 1.1 holds without restrictions such as condition (1.3). Indeed, suppose that $\mathbb{E}\{\lambda_d(X(\mathbb{R}_+^N))\} > 0$. Then, by Theorem 2.1, there exists a nonrandom $t \in \mathbb{R}_+^N$, and a $\mu \in \mathcal{P}([0, t])$ such that $\|\mu\|_e < +\infty$. Proposition A.3, then, shows that $\|\lambda_N \lfloor_{[0, t]}\|_e < +\infty$. We now appeal to Theorem 1.3 to deduce the existence of a local time process, $\mathbb{R}_+^N \ni t \mapsto \ell_t(\cdot) = L_{\lambda_N \lfloor_{[0, t]}}(\cdot)$, such that for all $t \in \mathbb{R}_+^N$, and all bounded measurable $f : \mathbb{R}^d \rightarrow \mathbb{R}$, the following holds \mathbb{P} -a.s.:

$$\int_{[0, t]} f(X(s)) ds = \int_{\mathbb{R}^d} f(a) \ell_t(a) da.$$

Moreover, $\ell_t \in L^2(\mathbb{R}^d)$, \mathbb{P} -almost surely. Theorem 1.1 can be shown to follow, if we could show that this fact would imply that $\int_{\mathbb{R}_+^N} \exp(-\sum_{j=1}^N t_j) \|\ell_t\|_{L^2(\mathbb{R}^d)} dt$ has a finite expectation. When $N = 1$, this follows from the strong Markov property. [Of course, when $N = 1$, condition (1.3) holds tautologically.] However, when $N > 1$, we do not know if such a fact holds.

Acknowledgment. The authors thank the referee for his or her careful reading of the paper and for several thoughtful comments that led to several improvements of the manuscript.

REFERENCES

- ADLER, R. J. (1981). *The Geometry of Random Fields*. Wiley, New York.
- BAKRY, D. (1979). Sur la régularité des trajectoires des martingales deux indices. *Z. Wahrsch. Verw. Gebiete* **50** 149–157.
- BENDIKOV, A. (1994). Asymptotic formulas for symmetric stable semigroups. *Expo. Math.* **12** 381–384.
- BERTOIN, J. (1996). *Lévy Processes*. Cambridge Univ. Press.
- BLUMENTHAL, R. M. and GETTOOR, R. (1961). Sample functions of stochastic processes with stationary independent increments. *J. Math. Mech.* **10** 493–516.
- BRETAGNOLLE, J. (1971). Résultats de Kesten sur les processus à accroissements indépendants. *Séminaire de Probabilités V. Lecture Notes in Math.* **191** 21–36. Springer, Berlin.
- CAIROLI, R. (1970). Une inégalité pour martingales à indices multiples et ses applications. *Séminaire de Probabilités VI. Lecture Notes in Math.* **124** 1–27. Springer, Berlin.
- CARLESON, L. (1983). *Selected Problems on Exceptional Sets*. Wadsworth, Belmont, CA.
- DELLACHERIE, C. and MEYER, P.-A. (1978). *Probabilities and Potential*. North-Holland, Amsterdam.
- EHM, W. (1981). Sample function properties of multi-parameter stable processes. *Z. Wahrsch. Verw. Gebiete* **56** 195–228.
- EVANS, S. N. (1987a). Multiple points in the sample paths of a Lévy process. *Probab. Theory Related Fields* **76** 359–367.
- EVANS, S. N. (1987b). Potential theory for a family of several Markov processes. *Ann. Inst. H. Poincaré Probab. Statist.* **23** 499–530.
- FITZSIMMONS, P. J. and SALISBURY, T. S. (1989). Capacity and energy for multiparameter processes. *Ann. Inst. H. Poincaré Probab. Statist.* **25** 325–350.
- FRISTEDT, B. (1974). Sample functions of stochastic processes with stationary, independent increments. *Advances in Probability and Related Topics* **3** 241–396.
- GEMAN, D. and HOROWITZ, J. (1980). Occupation densities. *Ann. Probab.* **8** 1–67.
- HAWKES, J. (1971). A lower Lipschitz condition for the stable subordinator. *Z. Wahrsch. Verw. Gebiete* **17** 23–32.
- HAWKES, J. (1974). Local times and zero sets for processes with infinitely divisible distributions. *J. London Math. Soc.* **8** 517–525.
- HAWKES, J. (1979). Potential theory of Lévy processes. *Proc. London Math. Soc.* **38** 335–352.
- HAWKES, J. (1986). Local times as stationary processes. In *From Local Times to Global Geometry* (K. D. Ellworthy, ed.) 111–120. Longman, Chicago.
- HENDRICKS, W. J. (1983). A uniform lower bound for Hausdorff dimension for transient symmetric Lévy processes. *Ann. Probab.* **11** 589–592.
- HIRSCH, F. (1995). Potential theory related to some multiparameter processes. *Potential Anal.* **4** 245–267.

- HIRSCH, F. and SONG, S. (1995a). Symmetric Skorohod topology on n -variable functions and hierarchical Markov properties of n -parameter processes. *Probab. Theory Related Fields* **103** 25–43.
- HIRSCH, F. and SONG, S. (1995b). Markov properties of multiparameter processes and capacities. *Probab. Theory Related Fields* **103** 45–71.
- HU, X. (1994). Some fractal sets determined by stable processes. *Probab. Theory Related Fields* **100** 205–225.
- KAHANE, J.-P. (1985). *Some Random Series of Functions*, 2nd ed. Cambridge Univ. Press.
- KESTEN, H. (1969). *Hitting Probabilities of Single Points for Processes with Stationary Independent Increments*. Amer. Math. Soc., Providence RI.
- KHOSHNEVISAN, D. and XIAO, Y. (2000). Level sets of additive random walks. In *High Dimensional Probability* (E. Giné, D. M. Mason and J. A. Wellner, eds.) 329–345. Birkhäuser, Basel.
- KHOSHNEVISAN, D. and XIAO, Y. (2002a). Level sets of additive Lévy processes. *Ann. Probab.* **30** 62–100.
- KHOSHNEVISAN, D. and XIAO, Y. (2002b). Weak unimodality of finite measures, and an application to potential theory of additive Lévy processes. *Proc. Amer. Math. Soc.* To appear.
- KHOSHNEVISAN, D., XIAO, Y. and ZHONG, Y. (2002). Local times of additive Lévy processes: Regularity. *Stochastic Process. Appl.* To appear.
- OREY, S. (1967). Polar sets for processes with stationary independent increments. In *Markov Processes and Potential Theory* (J. Chover, ed.) 117–126. Wiley, New York.
- PRUITT, W. E. (1969). The Hausdorff dimension of the range of a process with stationary independent increments. *J. Math. Mech.* **19** 371–378.
- PRUITT, W. E. and TAYLOR, S. J. (1996). Packing and covering indices for a general Lévy process. *Ann. Probab.* **24** 971–986.
- ROZANOV, YU. A. (1982). *Markov Random Fields*. Springer, Berlin.
- TAYLOR, S. J. (1973). Sample path properties of processes with stationary independent increments. In *Stochastic Analysis (A Tribute to the Memory of Rollo Davidson)* 387–414. Wiley, London.
- WALSH, J. B. (1986). Martingales with a multidimensional parameter and stochastic integrals in the plane. *Lecture Notes in Math.* **1215** 329–491. Springer, Berlin.

D. KHOSHNEVISAN
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF UTAH
JWB 233
155 S. 1400 E.
SALT LAKE CITY, UTAH 84112-0090
E-MAIL: davar@math.utah.edu
URL: <http://www.math.utah.edu/~davar>

Y. XIAO
DEPARTMENT OF STATISTICS
AND PROBABILITY
A-413 WELLS HALL
MICHIGAN STATE UNIVERSITY
EAST LANSING, MICHIGAN 48824
E-MAIL: xiao@stt.msu.edu
URL: <http://www.stt.msu.edu/~xiaoyimi>

Y. ZHONG
INSTITUTE OF APPLIED MATHEMATICS
ACADEMIA SINICA
BEIJING, 100080
PEOPLE'S REPUBLIC OF CHINA