



Reprint of: Boundary conditions for fractional diffusion[☆]

Boris Baeumer^a, Mihály Kovács^b, Mark M. Meerschaert^{c,*},
Harish Sankaranarayanan^d

^a University of Otago, New Zealand

^b Chalmers University of Technology and the University of Gothenburg, Sweden

^c Department of Statistics and Probability, Michigan State University, USA

^d Michigan State University, USA



ARTICLE INFO

Article history:

Received 26 May 2017

Received in revised form 18 December 2017

Keywords:

Fractional calculus

Boundary value problem

Numerical solution

Well-posed

ABSTRACT

This paper derives physically meaningful boundary conditions for fractional diffusion equations, using a mass balance approach. Numerical solutions are presented, and theoretical properties are reviewed, including well-posedness and steady state solutions. Absorbing and reflecting boundary conditions are considered, and illustrated through several examples. Reflecting boundary conditions involve fractional derivatives. The Caputo fractional derivative is shown to be unsuitable for modeling fractional diffusion, since the resulting boundary value problem is not positivity preserving.

© 2018 Published by Elsevier B.V.

1. Introduction

The space-fractional diffusion equation replaces the second derivative or Laplacian in the traditional diffusion equation with a fractional derivative. Fractional derivatives were invented soon after their integer-order counterparts, and by now have become an established field of study with a wide variety of applications in science and technology [1–8]. Practical applications include physics [9–12], finance [13–17], microbiology [18–20], medical imaging [21–26], and hydrology [27–30]. Zaslavsky [31,32] initiated the application of fractional calculus to chaotic dynamical systems. These methods may also be applicable to nonlinear dynamics in Hodgkin–Huxley neurons and pancreatic beta cells [33–35]. Many effective numerical methods have been developed for fractional differential equations, along with proofs of stability and consistency [7,36–50]. Because fractional derivatives are nonlocal operators, the concept of a boundary condition takes on new meaning [51,52], and the specification, implementation, and interpretation of physically meaningful boundary conditions remains an open problem.

Meerschaert and Tadjeran [42,43] consider a fractional diffusion equation with Dirichlet boundary conditions, but the physical meaning of those boundary conditions was not addressed. Jin et al. [53] establish variational formulations for fractional differential equations with zero Dirichlet boundary conditions. Ferreira [54] establishes a Lyapunov inequality for a fractional equation with zero Dirichlet boundary conditions. Zhu et al. [55] prove a Lyapunov formula for a fractional equation with periodic boundary conditions. D. del-Castillo-Negrete [56] considers space–time fractional diffusions on an interval with absorbing boundary conditions. Montefusco et al. [57] consider a fractional reaction–diffusion equation with a traditional Neumann boundary condition. Xie et al. [58] consider a fractional Poisson equation with traditional Neumann

DOI of original article: <http://dx.doi.org/10.1016/j.cam.2017.12.053>.

[☆] This article is a reprint of a previously published article [B. Baeumer et al. J. Comput. Appl. Math. 336 (2018) 408–424].

* Corresponding author.

E-mail addresses: bbaeumer@maths.otago.ac.nz (B. Baeumer), mihaly@chalmers.se (M. Kovács), mcubed@stt.msu.edu (M.M. Meerschaert), harish@msu.edu (H. Sankaranarayanan).

<https://doi.org/10.1016/j.cam.2018.03.007>

0377-0427/© 2018 Published by Elsevier B.V.

boundary conditions. Laskin [59] solves a fractional Schrödinger equation with reflecting boundary conditions using a square potential. Szekeres and Izsák [60] solve fractional diffusion equations with zero Dirichlet and traditional Neumann boundary conditions via periodic extensions. Lim and Teo [61,62] apply both the traditional Neumann condition and a fractional Neumann condition to study the finite temperature Casimir force acting on a piston moving freely inside a rectangular cavity. Wang and Yang [63] consider a fractional Poisson equation with both traditional and fractional Neumann boundary conditions, and they show that the appropriate choice of boundary condition depends on the type of fractional derivative. Voller [64] solves a fractional Stefan problem, with a nonzero Dirichlet condition on the left, and a moving boundary on the right. Kolokoltsov [65] implements Dirichlet and Neumann conditions by modifying the jump intensity in the Lévy representation. Warma [66] gives an abstract characterization of Neumann boundary conditions for the fractional Laplacian based on Dirichlet forms. Umarov [67] proves existence and uniqueness for general nonlocal diffusions with Wentzel's boundary conditions, defined by a pseudo-differential operator. Du et al. [68] develop Dirichlet and Neumann volume constraints based on the theory of nonlocal diffusion. Dipierro et al. propose a new kind of Neumann condition that restricts integration in the definition of the fractional Laplacian to the boundary. Krepsheva et al. [69] implement an elastic reflecting boundary condition for the fractional Laplacian by modifying the integration kernel. Cusimano et al. [70] and Ilic et al. [71] take a different approach to elastic reflection, taking limits of a discrete fractional Laplacian.

This paper considers space-fractional diffusion equations on the unit interval $0 \leq x \leq 1$ with absorbing or reflecting boundary conditions. Both Riemann–Liouville and Caputo flux forms are considered, and the profound difference in their solutions is illustrated. To specify a fractional diffusion equation on a bounded domain, appropriate boundary conditions must be enforced [51,52]. We discuss absorbing (Dirichlet) and reflecting (Neumann) boundary conditions, which can take a very different form for a fractional evolution equation. In particular, the appropriate Neumann boundary condition sets a fractional derivative equal to zero at the boundary, not the first derivative as in the traditional diffusion equation. We also show that the Caputo form does not preserve positivity, and hence cannot provide a suitable model for anomalous diffusion.

By varying the type of space-fractional derivative and the boundary conditions, we obtain a number of possible fractional diffusion equations on the unit interval. For each of these, we develop and apply a suitable numerical solution method. We demonstrate that fractional Neumann boundary conditions are physically meaningful zero-flux conditions that correspond to inelastic reflection. We also show that the traditional Neumann condition is not satisfied at the boundary. We then review the underlying theory from the point of view of abstract evolution equations, semigroups and generators. Well-posedness is verified, including uniqueness of solutions. Steady-state solutions are identified, and convergence to steady state is demonstrated.

2. Fractional boundary value problems

Consider the fractional diffusion equation

$$\frac{\partial}{\partial t} u(x, t) = C \mathbb{D}^\alpha u(x, t) \quad \text{for } 1 < \alpha < 2 \quad (2.1)$$

on the entire real line, where the Riemann–Liouville fractional derivative

$$\mathbb{D}^\alpha u(x, t) = \frac{1}{\Gamma(n - \alpha)} \frac{\partial^n}{\partial x^n} \int_{-\infty}^x u(y, t) (x - y)^{n - \alpha - 1} dy \quad (2.2)$$

for $\alpha > 0$ and $n - 1 < \alpha \leq n$. Note that (2.2) is a nonlocal operator that depends on the values of $u(y, t)$ at every point $y < x$. The exact analytical solution to (2.1) can be written in terms of a stable probability density function. Although this analytical solution cannot be computed in closed form, there are readily available codes that compute the stable density, and these can be used to plot the solutions to (2.1). See for example [72, Chapter 5].

However, if we restrict the fractional diffusion to a finite interval, then there are no known analytical solutions, and numerical methods must be used. First consider the fractional diffusion equation

$$\frac{\partial}{\partial t} u(x, t) = C \mathbb{D}_{[0,x]}^\alpha u(x, t) \quad \text{for } 1 < \alpha < 2 \quad (2.3)$$

on the state space $0 \leq x \leq 1$ with initial condition $u(x, 0) = u_0(x) \geq 0$. On this finite domain, we define the Riemann–Liouville fractional derivative

$$\mathbb{D}_{[0,x]}^\alpha u(x, t) = \frac{1}{\Gamma(n - \alpha)} \frac{\partial^n}{\partial x^n} \int_0^x u(y, t) (x - y)^{n - \alpha - 1} dy, \quad (2.4)$$

the only difference from (2.2) being the lower limit of integration. This is still a nonlocal operator, since it depends on the values of $u(y, t)$ at every point $0 \leq y \leq x$.

3. Absorbing boundary conditions

Now let us impose a zero boundary condition at each endpoint:

$$u(0, t) = u(1, t) = 0 \quad \text{for all } t \geq 0. \quad (3.1)$$

The zero Dirichlet boundary conditions (3.1) are usually called absorbing boundary conditions, but do they have the same meaning for a nonlocal operator? To illuminate this issue, let us develop a numerical method to solve the fractional diffusion equation (2.3), paying special attention to meaning of the zero boundary conditions (3.1).

The fractional derivative (2.2) can be approximated using the Grünwald–Letnikov formula [72, Proposition 2.1]

$$\mathbb{D}^\alpha u(x, t) = \lim_{h \rightarrow 0} h^{-\alpha} \sum_{i=0}^{\infty} (-1)^i \binom{\alpha}{i} u(x - ih, t) \tag{3.2}$$

where the Grünwald weights are given by

$$g_i^\alpha = (-1)^i \binom{\alpha}{i} = \frac{(-1)^i \Gamma(\alpha + 1)}{\Gamma(i + 1) \Gamma(\alpha - i + 1)} \tag{3.3}$$

for all $i \geq 0$. Since the finite domain fractional derivative (2.4) is equivalent to the Riemann–Liouville fractional derivative of a function that vanishes for $x < 0$, we immediately obtain that

$$\mathbb{D}_{[0,x]}^\alpha u(x, t) = \lim_{h \rightarrow 0} h^{-\alpha} \sum_{i=0}^{\lfloor x/h \rfloor} g_i^\alpha u(x - ih, t). \tag{3.4}$$

This approximation can be used to construct numerical solutions to the fractional diffusion equation, but the resulting methods are unstable [42, Proposition 2.3]. Instead, we apply a shifted Grünwald formula

$$\mathbb{D}_{[0,x]}^\alpha u(x, t) \approx h^{-\alpha} \sum_{i=0}^{\lfloor x/h \rfloor + 1} g_i^\alpha u(x - (i - 1)h, t) \tag{3.5}$$

which results in a stable method [42, Theorem 2.7].

To illuminate the role of the boundary conditions, first consider the fractional diffusion equation (2.1) on the real line. As a thought experiment, discretize $x_j = jh$ and $t_k = k\Delta t$ and apply the Grünwald approximation to obtain the explicit Euler scheme

$$u(x_j, t_{k+1}) = u(x_j, t_k) + Ch^{-\alpha} \sum_{i=0}^{\infty} g_i^\alpha u(x_{j-i+1}, t_k) \Delta t. \tag{3.6}$$

The Grünwald weights are $g_0^\alpha = 1, g_1^\alpha = -\alpha, g_2^\alpha = \alpha(\alpha - 1)/2!$ and so forth, and note that $g_i^\alpha > 0$ for all $i \neq 1$. The scheme is mass-preserving because [72, Eq. (2.11)]

$$\sum_{i=0}^{\infty} g_i^\alpha = 0, \tag{3.7}$$

and hence to understand a physical model of the fractional diffusion, it will suffice to consider $u(x_j, t_k)h$ as the mass at location x_j at time t_k . The total mass $M_k = \sum_j u(x_j, t_k)h$ does not vary with time t_k , but rather remains equal to the initial mass $M_0 = \sum_j u_0(x_j)h$. The scheme moves a mass $C\Delta t h^{-\alpha-1} g_i^\alpha u(x_{j-i+1}, t_k)h$ from location x_{j-i+1} to location x_j when $i \neq 1$. The total mass $C\Delta t h^{-\alpha} \alpha u(x_j, t_k)$ moved out of location x_j is equal to the sum of the amounts moved from location x_j to another location, because $\sum_{i \neq 1} g_i^\alpha = \alpha$. In this scheme, mass can be transported large distances to the right, but only one step size h to the left. Note that the scheme (3.6) is also positivity preserving for $C\alpha h^{-\alpha} \Delta t \leq 1$, since a fraction $\leq 100\%$ of the mass at each point is removed, and then redistributed.

Now we want to restrict to the unit interval $0 \leq x \leq 1$ and impose the zero boundary conditions (3.1). Since we are solving a nonlocal problem, this requires some care. Unlike a traditional diffusion equation, the Euler scheme (3.6) moves mass a long distance in one time step, for any step size. That mass can land outside the unit interval, and then it must be accounted for in the scheme. Part of the picture is to understand how the Grünwald approximation (3.4) accounts for this mass. The remaining part is to understand the zero boundary conditions.

Let us note that the discretization of the fractional diffusion equation (2.3) on the bounded domain using (3.5) takes the form

$$u(x_j, t_{k+1}) = u(x_j, t_k) + Ch^{-\alpha} \sum_{i=0}^{j+1} g_i^\alpha u(x_{j-i+1}, t_k) \Delta t, \quad \forall 0 \leq j \leq n. \tag{3.8}$$

Comparing with (3.6), we can see that no mass is moved to location x_j from any location x_{j-i+1} when $i > j + 1$, i.e., when $x_{j-i+1} < 0$ lies outside the domain $0 \leq x \leq 1$.

Now we impose the boundary conditions (3.1) by setting $u(x_j, t_k) = 0$ when $j = 0$ (location $x_j = 0$) or $j = n$ (location $x_j = 1$), where $nh = 1$. Since our initial condition $u_0(x)$ must also satisfy the boundary conditions, we start with all the mass

inside the open interval $0 < x < 1$. To enforce the boundary conditions, we have to modify the Euler scheme (3.8). After a simple change of variables, we can write (3.8) in the form

$$u(x_j, t_{k+1}) = u(x_j, t_k) + Ch^{-\alpha} \sum_{i=0}^n b_{ij} u(x_i, t_k) \Delta t, \quad \forall 0 \leq j \leq n, \tag{3.9}$$

where $b_{ij} = g_{j-i+1}^\alpha$ for $i \leq j + 1$ and $b_{ij} = 0$ for $i > j + 1$. Next we will modify certain coefficients b_{ij} to enforce the boundary conditions. First consider the left end point $x_0 = 0$. Since the mass at this location has to remain zero,

$$0 = \sum_{i=0}^1 b_{i0} u(x_i, t_k).$$

Since $u(x_0, t_k) = 0$ for all k , this requires $b_{10} = 0$. Now the mass $C \Delta t h^{-\alpha-1} g_0^\alpha u(x_1, t_k) h$ that would have been transported from location x_1 to location x_0 is instead removed from the system, to enforce the zero boundary condition. Next consider the right end point $x_n = 1$. Since the mass at this location has to remain zero, we require

$$0 = \sum_{i=0}^n b_{in} u(x_i, t_k).$$

Since $u(x_n, t_k) = 0$ for all k , and since all $u(x_i, t_k) \geq 0$ for step size $\Delta t \leq h^\alpha / C\alpha$ and a nonnegative initial condition, we must have $b_{in} = 0$ for all $i = 0, 1, 2, n - 1$. This change alters (3.8) by taking the mass $C \Delta t h^{-\alpha} g_{n-i+1}^\alpha u(x_i, t_k)$ that would have been transported from location $x_i < 1$ to location $x_n = 1$ and removing it from the system. The resulting scheme can be written in the form (3.9) where

$$b_{ij} = \begin{cases} g_{j-i+1}^\alpha & \text{if } 0 < j < n \text{ and } i \leq j + 1, \\ 0 & \text{otherwise.} \end{cases} \tag{3.10}$$

To interpret (3.10), recall that $Ch^{-\alpha} b_{ij} u(x_i, t_k) \Delta t$ is the mass transferred from location x_i to location x_j during this time step.

In summary, the fractional diffusion equation (2.3) on $0 \leq x \leq 1$ with zero boundary conditions (3.1) is indeed a model with absorbing boundary conditions. As compared to the Euler scheme on the entire real line, here the mass scheduled for transport to or beyond the boundary of the unit interval is instead deleted from the system, or absorbed. This scheme is also positivity preserving so long as $C\alpha \Delta t h^{-\alpha} \leq 1$, since a fraction of the mass at each point is removed, and then redistributed or absorbed.

Write $\beta = Ch^{-\alpha} \Delta t$, $u_j^k = u(x_j, t_k)$, the solution vector $\mathbf{u}_k = [u_0^k, \dots, u_n^k]$, and the $(n+1) \times (n+1)$ iteration matrix $B = [b_{ij}]$. Then we can express the explicit Euler scheme (3.9) in vector-matrix form

$$\mathbf{u}_{k+1} = \mathbf{u}_k + \beta \mathbf{u}_k B. \tag{3.11}$$

In this form, the ij entry of the matrix B is proportional to the rate at which mass is transferred from location x_i to location x_j . Equivalently, we can write

$$\mathbf{u}_{k+1}^T = \mathbf{u}_k^T + \beta B^T \mathbf{u}_k^T. \tag{3.12}$$

The formulation (3.12) is traditional in numerical analysis, e.g., see [43, p. 4], while (3.11) is used for Markov chains, e.g., see [73, Section 8.1].

The explicit Euler scheme (3.12) is stable under a step size condition $\alpha\beta \leq 1$, or equivalently, $C\alpha \Delta t \leq h^\alpha$, see [43, Proposition 2.1]. The implicit Euler scheme

$$\mathbf{u}_{k+1}^T = \mathbf{u}_k^T + \beta B^T \mathbf{u}_{k+1}^T \tag{3.13}$$

is unconditionally stable [42, Theorem 2.7]. As noted in the Introduction, by now there are a wide variety of numerical methods to solve this problem. For example, the explicit Euler scheme (3.11) can be viewed as the temporal discretization of a linear system of ordinary differential equations (method of lines, e.g., see [40,74]), and then any standard method for solving the linear system can be employed.

Fig. 1 shows a numerical solution of the fractional diffusion equation (2.3) on $0 \leq x \leq 1$ with zero boundary conditions (3.1). The solution was plotted using the MATLAB routine `ode15s` for stiff systems of ordinary differential equations, viewing the explicit Euler scheme (3.11) as the temporal discretization of a linear system of ordinary differential equations (method of lines), with time step $\Delta t = 0.01$ and spatial grid size $h = 0.001$. The tent function initial condition

$$u_0(x) = \begin{cases} 25x - 7.5 & \text{for } 0.3 < x \leq 0.5, \\ -25x + 17.5, & \text{for } 0.5 < x < 0.7, \\ 0 & \text{otherwise} \end{cases} \tag{3.14}$$




Fig. 1. Numerical solution of the fractional diffusion equation (2.3) with $\alpha = 1.5$ and $C = 1$ on $0 \leq x \leq 1$ with zero boundary conditions at time $t = 0$ (solid line), $t = 0.05$ (dashed), $t = 0.1$ (dash dot), $t = 0.5$ (dotted).

satisfies the zero boundary conditions, and integrates to total mass $M = 1$. Because of the absorbing boundary conditions, solutions tend to zero as t increases. Refining the temporal or spatial discretization resulted in no visible change in the plots. Because the fractional derivative (2.4) is one sided, solution curves are skewed for all $t > 0$, even though the initial mass distribution is symmetric. This can also be seen from (3.8), since the mass $\beta g_0^\alpha u_i^k$ moved from state x_i to state x_{i-1} exceeds the total amount of mass moved to the right (which is less than $\beta(\alpha - 1)u_i^k$), at any node inside the domain. We have also written explicit Euler codes for all the examples in this paper using the free and open source R programming language [75]. Please contact the authors to obtain the MATLAB or R codes.

Remark 3.1. A few crucial differences from the traditional diffusion setup should be noted. First of all, one can also characterize the physical problem as absorbing on the exterior of the open domain $0 < x < 1$, not just at the boundary. Physically, mass can be displaced a long distance from the domain, and then absorbed. Second, the form of the fractional derivative (2.4) also incorporates absorbing outside the domain. The fractional diffusion equation (2.1) on the real line with the exterior condition $u(x, t) = 0$ for $x \leq 0$ or $x \geq 1$ is equivalent to the fractional diffusion equation (2.3) on the bounded domain $0 \leq x \leq 1$ with zero boundary conditions (3.1). The fractional derivative itself codes the zero exterior condition on $x < 0$. For more details, and an interesting connection to stochastic processes, see [51]. Third, since the positive Riemann–Liouville fractional derivative (2.2) is one-sided, depending only on values of the function to the left, the zero exterior condition on $x \geq 1$ is automatically enforced. Another way to see this is that, in the Euler scheme, mass can be transported to location x_j from any location to the left, but not from the right.

Remark 3.2. Theoretical properties of the solution are discussed in [51,76]. There it is shown that the Cauchy problem (2.3) on $0 \leq x \leq 1$ with zero boundary conditions (3.1) (or equivalently, zero exterior condition) is well-posed: There exists a unique solution for any initial condition $u_0(x)$ that depends continuously on this initial function. The general theory in [51] applies on the Banach space $C_0(0, 1)$ of continuous functions that vanish at the end points, with the supremum norm. In [76] the Banach space $L^1[0, 1]$ is considered. Since both the implicit and explicit Euler methods are consistent, and stable (in the explicit case, under a step size condition on Δt), and since the problem (2.3) on $0 \leq x \leq 1$ with zero boundary conditions is well-posed, the Lax Equivalence Theorem [77, p. 45] implies that either of these Euler methods will converge to the unique solution as $h \rightarrow 0$ and $\Delta t \rightarrow 0$. The same is true for any other stable, consistent numerical method. The theory in [51] also relates the Cauchy problem (2.3) on $0 \leq x \leq 1$ with zero boundary conditions to a probability model, which implies that the problem is positivity preserving. The analysis in [76] also computes the exact domain of the generator $\mathbb{D}_{[0,x]}^\alpha$ on $L^1[0, 1]$ with zero boundary conditions.

Remark 3.3. The astute reader will notice that (3.8) with $j = n$ involves the mass at location $x_{n+1} = 1 + h$ when $j = n$, and this x_{n+1} term does not appear in (3.9). We could indeed track the mass moved to the location x_{n+1} , which is outside the domain, but with the zero boundary condition $u(x_n, t_k) = 0$, none of this mass can ever come back into the domain. Indeed, mass from location x_{n+1} can only move left one step to location $x_n = 1$, and the zero boundary condition forbids this. In other words, if we did include state x_{n+1} in our scheme, then we would also conclude $b_{n+1,n} = 0$ by the same argument that $b_{in} = 0$ for $0 \leq i \leq n - 1$. Hence we need not track the mass at this location.

4. Reflecting boundary conditions

The proper formulation of physically meaningful reflecting boundary conditions for the fractional diffusion equation (2.3) requires careful consideration of the nonlocal operator (2.4). Suppose that our goal is for mass leaving the domain to instead come to rest at the boundary. Unlike the traditional diffusion setup, this mass can come from far inside the domain, not just an adjacent grid point. Now the mass that was removed from the system in the Dirichlet model of Section 3 will instead be preserved, and moved to the boundary.

Let us consider the right boundary $x_n = 1$, since long movements are always to the right in our setup. For each $i = 1, 2, \dots, n - 1$, at each time step, mass $\beta\alpha u_i^k$ is moved out of location x_i , and redistributed. Of this total, a fraction $\beta g_{j-i+1}^\alpha u_i^k$ is moved to location x_j when $j = i - 1$ or $j > i$. Hence the mass landing at, or exiting the domain through, the right boundary $x_n = 1$ from location x_i for $i = 1, 2, \dots, n - 1$ is

$$\sum_{j=n}^{\infty} \beta g_{j-i+1}^\alpha u_i^k.$$

Then using (3.7), along with the identity [8, Eq. (20.4)]

$$\sum_{j=0}^n g_j^\alpha = g_n^{\alpha-1}, \tag{4.1}$$

we set $b_{in} = -g_{n-i}^{\alpha-1}$ in (3.9), for each $i = 1, 2, \dots, n - 1$. In the scheme (3.6) on the real line, the mass $\beta g_0^\alpha u_n^k$ moves from location x_n to location x_{n-1} , and the remainder of the mass $\beta\alpha u_n^k$ leaving location x_n moves to the right, outside the domain $0 \leq x \leq 1$. In the reflecting scheme, we retain this mass at location x_n by setting $b_{nn} = -1 = -g_0^\alpha$.

The only way that mass can move to the left boundary $x_0 = 0$ in this scheme is from the adjacent node $x_1 = h$, hence we leave $b_{10} = g_0^\alpha = 1$. In the scheme (3.6) on the real line, mass $\beta g_0^\alpha u_0^k$ moves from location x_0 to location $x_{-1} < 0$. To prevent this, and thus to keep the scheme mass-preserving, recall that $g_0^\alpha = 1$ and $g_1^\alpha = -\alpha$, and set $b_{00} = 1 - \alpha$. To prevent mass leaving state x_0 from jumping through the right boundary, we also set

$$b_{0n} = \sum_{j=n}^{\infty} g_{j+1}^\alpha = -g_n^{\alpha-1} > 0,$$

and hence the explicit Euler scheme for the case of reflecting boundary conditions is written in the form (3.9) with

$$b_{ij} = \begin{cases} g_{j-i+1}^\alpha & \text{if } 0 < j < n \text{ and } i \leq j + 1, \\ 1 & \text{if } i = 1 \text{ and } j = 0, \\ 1 - \alpha & \text{if } i = j = 0, \\ -g_{n-i}^{\alpha-1} & \text{if } j = n \text{ and } i \leq n, \\ 0 & \text{otherwise.} \end{cases} \tag{4.2}$$

Next we will argue that the reflecting boundary conditions for the fractional diffusion equation (2.3) on $0 \leq x \leq 1$ can be written in the form

$$\mathbb{D}_{[0,x]}^{\alpha-1} u(0, t) = \mathbb{D}_{[0,x]}^{\alpha-1} u(1, t) = 0 \quad \text{for all } t \geq 0, \tag{4.3}$$

using the Riemann–Liouville fractional derivative (2.4) of order $\alpha - 1$. When $\alpha = 2$, this reduces to the classical reflecting condition $\frac{\partial}{\partial x} u(x, t) = 0$ at the boundary. First consider the right boundary $x_n = 1$, and write out the iteration equation for this node: From (3.9) and (4.2) with $\beta = Ch^{-\alpha} \Delta t$ we have $u_n^{k+1} = u_n^k - \beta g_n^{\alpha-1} u_0^k - \dots - \beta g_1^{\alpha-1} u_{n-1}^k - \beta g_0^{\alpha-1} u_n^k$ which is algebraically equivalent to

$$h \frac{u_n^{k+1} - u_n^k}{\Delta t} = -Ch^{1-\alpha} \sum_{i=0}^n g_{n-i}^{\alpha-1} u_i^k = -Ch^{1-\alpha} \sum_{i=0}^n g_{n-i}^{\alpha-1} u(x_n - (n - i)h, t_k).$$

Letting $\Delta t \rightarrow 0$ and $h \rightarrow 0$, and using the Grünwald approximation (3.4), we arrive at the reflecting boundary condition (4.3) at the right boundary $x = 1$.

The iteration equation at the left boundary is $u_0^{k+1} = u_0^k + \beta(1 - \alpha)u_0^k + \beta u_1^k$. Recalling that $g_0^{\alpha-1} = 1$ and $g_1^{\alpha-1} = 1 - \alpha$, this reduces to

$$h \frac{u_0^{k+1} - u_0^k}{\Delta t} = Ch^{1-\alpha} \sum_{i=0}^1 g_{1-i}^{\alpha-1} u_i^k,$$

which is consistent with the reflecting boundary condition (4.3) at the left boundary $x = 0$. To rigorously prove that the left boundary condition in (4.3) holds, [76, Proposition 19] extends the matrix $h^{-\alpha} B$ by interpolation to an operator on $L^1[0, 1]$, and proves convergence to the generator (2.4) with boundary conditions (4.3).




Fig. 2. Numerical solution of the fractional diffusion equation (2.3) with $\alpha = 1.5$ and $C = 1$ on $0 \leq x \leq 1$ with reflecting boundary conditions (4.3) at time $t = 0$ (solid line), $t = 0.05$ (dashed), $t = 0.1$ (dash dot), $t = 0.5$ (dotted).

Remark 4.1. The reflecting boundary conditions (4.3) can be seen as zero flux conditions at the boundary: Note that the fractional diffusion equation (2.3) can be derived from the traditional conservation of mass equation

$$\frac{\partial}{\partial t} u(x, t) = -\frac{\partial}{\partial x} q(x, t) \tag{4.4}$$

together with the flux equation (or fractional Fick’s Law, see [78])

$$q(x, t) = -C \mathbb{D}_{[0,x]}^{\alpha-1} u(x, t) = -C \frac{\partial}{\partial x} \frac{1}{\Gamma(2-\alpha)} \int_0^x u(y, t)(x-y)^{1-\alpha} dy. \tag{4.5}$$

Hence (4.3) simply sets the flux to zero at the boundary. When $\alpha = 2$, the fractional Fick’s Law reduces to the traditional Fick’s Law $q(x, t) = -C \frac{\partial}{\partial x} u(x, t)$. D. del-Castillo-Negrete [56, Eq. (3)] noted that the flux in the Riemann–Liouville fractional diffusion equation is fractional.

Fig. 2 shows a numerical solution of the fractional diffusion equation (2.3) on $0 \leq x \leq 1$ with reflecting boundary conditions, using the same numerical method and initial function as in Fig. 1. As in Fig. 1, and for the same reason, solution curves are skewed for all $t > 0$, even though the initial mass distribution is symmetric. However, there is a profound difference in the solutions. Here the total mass (area under the curve) remains equal to the initial mass $M = 1$ for all $t > 0$, because of the reflecting boundary conditions. As t increases, the solutions approach the steady state solution $u_\infty(x) = (\alpha - 1)x^{\alpha-2}$ on $0 < x < 1$.

Remark 4.2. The general steady state solution to the fractional diffusion equation (2.3) is $u_\infty(x) = c_1 x^{\alpha-1} + c_2 x^{\alpha-2}$ where c_1, c_2 are arbitrary real numbers. To see this, note that the Riemann–Liouville fractional derivative $\mathbb{D}_{[0,x]}^\alpha u(x) = \frac{d^2}{dx^2} \mathbb{J}_{[0,x]}^{2-\alpha} u(x)$ where the Riemann–Liouville fractional integral

$$\mathbb{J}_{[0,x]}^\gamma u(x) = \frac{1}{\Gamma(\gamma)} \int_0^x u(y)(x-y)^{\gamma-1} dy \tag{4.6}$$

for any $\gamma > 0$. Using the general formula (e.g., see [72, Example 2.7])

$$\mathbb{J}_{[0,x]}^\gamma [x^p] = \frac{\Gamma(p+1)}{\Gamma(p+\gamma+1)} x^{p+\gamma} \tag{4.7}$$

we see that

$$\mathbb{J}_{[0,x]}^{2-\alpha} u_\infty(x) = c_1 \Gamma(\alpha) x + c_2 \Gamma(\alpha - 1).$$

Then $\mathbb{D}_{[0,x]}^\alpha u_\infty(x) = 0$ for all $0 < x < 1$. The only steady state solution with total mass 1 that satisfies the reflecting boundary conditions (4.3) has $c_1 = 0$ and $c_2 = \alpha - 1$. The only steady state solution that satisfies the absorbing boundary conditions (4.3) has $c_1 = 0$ and $c_2 = 0$.




Fig. 3. Numerical solution of the fractional diffusion equation (2.3) with $\alpha = 1.5$ and $C = 1$ on $0 \leq x \leq 1$ with boundary conditions (5.1): Reflecting on the left, and absorbing on the right at time $t = 0$ (solid line), $t = 0.05$ (dashed), $t = 0.1$ (dash dot), $t = 0.5$ (dotted).

5. Absorbing on one side, reflecting on the other

Next we consider the fractional diffusion equation (2.3) on $0 \leq x \leq 1$ with a reflecting boundary condition on the left, and an absorbing boundary condition on the right:

$$\mathbb{D}_{[0,x]}^{\alpha-1}u(0, t) = 0 \quad \text{and} \quad u(1, t) = 0 \quad \text{for all } t \geq 0 \tag{5.1}$$

The explicit Euler scheme for this problem is (3.9) with

$$b_{ij} = \begin{cases} g_{j-i+1}^\alpha & \text{if } 0 < j < n \text{ and } i \leq j + 1, \\ 1 & \text{if } i = 1 \text{ and } j = 0, \\ 1 - \alpha & \text{if } i = j = 0, \\ 0 & \text{otherwise.} \end{cases} \tag{5.2}$$

This combines the reflecting boundary condition at $x_0 = 0$ from (4.2) and the absorbing boundary condition at $x_n = 1$ from (3.10).

Fig. 3 shows the resulting numerical solution of the fractional diffusion equation (2.3) on $0 \leq x \leq 1$ with boundary conditions (5.1), using the same numerical method and initial function as in Fig. 1. The solutions are skewed to the right, and approach the steady state solution $u_\infty = 0$ as t increases. In this model, mass accumulates at the reflecting boundary $x = 0$, but then will eventually be absorbed at the right boundary $x = 1$.

Next we consider the opposite case, the fractional diffusion equation (2.3) on $0 \leq x \leq 1$ with an absorbing boundary condition on the left, and a reflecting boundary condition on the right:

$$u(0, t) = 0 \quad \text{and} \quad \mathbb{D}_{[0,x]}^{\alpha-1}u(1, t) = 0 \quad \text{for all } t \geq 0. \tag{5.3}$$

The explicit Euler scheme for this problem is (3.9) with

$$b_{ij} = \begin{cases} g_{j-i+1}^\alpha & \text{if } 0 < j < n \text{ and } i \leq j + 1, \\ -g_{n-i}^{\alpha-1} & \text{if } j = n \text{ and } i \leq n, \\ 0 & \text{otherwise.} \end{cases} \tag{5.4}$$

This combines the absorbing boundary condition at $x_0 = 0$ from (3.10) and the reflecting boundary condition at $x_n = 1$ from (4.2).

Fig. 4 shows the resulting numerical solution of the fractional diffusion equation (2.3) on $0 \leq x \leq 1$ with boundary conditions (5.3), using the same numerical method and initial function as in Fig. 1. The solutions are skewed to the right, and approach the steady state solution $u_\infty = 0$ as t increases. In this model, mass is reflected at the right boundary, and then eventually absorbed at the left boundary.

Remark 5.1. In [76, Proposition 19] it is shown that the Cauchy problem (2.3) on $0 \leq x \leq 1$ with boundary conditions (4.3) or (5.1) or (5.3) is well-posed on the Banach space $L^1[0, 1]$, and the exact domain of the generator is computed. Then it follows as in Remark 3.2 that any stable and consistent numerical method converges to the unique solution.




Fig. 4. Numerical solution of the fractional diffusion equation (2.3) with $\alpha = 1.5$ and $C = 1$ on $0 \leq x \leq 1$ with boundary conditions (5.3): Reflecting on the right, and absorbing on the left at time $t = 0$ (solid line), $t = 0.05$ (dashed), $t = 0.1$ (dash dot), $t = 0.5$ (dotted).

6. Caputo fractional flux

An alternative to the fractional diffusion equation (2.3) is the Caputo fractional flux model. The Caputo fractional derivative is defined by

$$\partial_{[0,x]}^\gamma u(x) = \frac{1}{\Gamma(n - \gamma)} \int_0^x u^{(n)}(y)(x - y)^{n-\gamma-1} dy \tag{6.1}$$

for $\gamma > 0$ and $n - 1 < \gamma \leq n$, where $u^{(n)}(x)$ is the n th derivative. It differs from (2.4) in that the derivative is moved inside the integral. These two fractional derivatives are not equivalent. For example,

$$\partial_{[0,x]}^\gamma u(x) = \mathbb{D}_{[0,x]}^\gamma u(x) - u(0) \frac{x^{-\gamma}}{\Gamma(1 - \gamma)} \tag{6.2}$$

when $0 < \gamma < 1$ [72, Eq. (2.33)]. Recall from Remark 4.1 that the fractional diffusion equation (2.3) can be derived from the conservation of mass equation (4.4) and the Riemann–Liouville fractional Fick’s Law (4.5). Noting that Fick’s Law is purely empirical, we can instead consider the Caputo fractional flux

$$q(x, t) = -C \partial_{[0,x]}^{\alpha-1} u(x, t) = -\frac{C}{\Gamma(2 - \alpha)} \int_0^x u'(y, t)(x - y)^{1-\alpha} dy, \tag{6.3}$$

where $u'(x, t)$ denotes the x derivative. This leads to the fractional diffusion equation with Caputo flux:

$$\frac{\partial}{\partial t} u(x, t) = C \mathbf{D}_{[0,x]}^\alpha u(x, t), \tag{6.4}$$

where the *Patie–Simon fractional derivative* is defined by

$$\mathbf{D}_{[0,x]}^\alpha u(x) = \frac{1}{\Gamma(2 - \alpha)} \frac{\partial}{\partial x} \int_0^x \frac{\partial}{\partial x} u(x - y)y^{-\alpha} dy \tag{6.5}$$

for $1 < \alpha < 2$. Patie and Simon [79, p. 570] showed that this operator is the (backward) generator of a standard spectrally negative α -stable process reflected to stay positive [79, p. 573]. Use the relation (6.2) and the definition (2.4) to see that

$$\begin{aligned} \mathbf{D}_{[0,x]}^\alpha f(x) &= \frac{d}{dx} [\partial_{[0,x]}^{\alpha-1} f(x)] \\ &= \frac{d}{dx} \left[\mathbb{D}_{[0,x]}^{\alpha-1} f(x) - f(0) \frac{x^{1-\alpha}}{\Gamma(2 - \alpha)} \right] \\ &= \mathbb{D}_{[0,x]}^\alpha f(x) - f(0) \frac{x^{-\alpha}}{\Gamma(1 - \alpha)} \end{aligned} \tag{6.6}$$

which relates the two derivatives when $1 < \alpha < 2$.

A Grünwald finite difference scheme for the fractional derivative (6.5) can be written as

$$\mathbb{D}_{[0,x]}^\alpha f(x) = \lim_{h \rightarrow 0} h^{-\alpha} \left[\sum_{i=0}^{j+1} g_i^\alpha f(x - (i-1)h) - g_{j+1}^{\alpha-1} f(x - jh) \right] \tag{6.7}$$

where $j = j(h) = \lfloor x/h \rfloor$. To see this, apply [72, Proposition 2.1] to see that the first term in (6.7) converges to $\mathbb{D}_{[0,x]}^\alpha f(x)$. Then note that $x - jh \rightarrow 0$ as $h \rightarrow 0$, and that [72, Eq. (2.5)]

$$g_j^{\alpha-1} \sim \frac{1-\alpha}{\Gamma(2-\alpha)} j^{-\alpha} \quad \text{as } j \rightarrow \infty, \tag{6.8}$$

meaning that the ratio between the left and right terms tends to 1 as $j \rightarrow \infty$. Then

$$\begin{aligned} h^{-\alpha} g_{j+1}^{\alpha-1} &\sim h^{-\alpha} \frac{1-\alpha}{\Gamma(2-\alpha)} ([x/h] + 1)^{-\alpha} \\ &\sim \frac{1-\alpha}{\Gamma(2-\alpha)} x^{-\alpha} = \frac{x^{-\alpha}}{\Gamma(1-\alpha)} \end{aligned}$$

using $\Gamma(z+1) = z\Gamma(z)$. Then (6.7) follows using (6.6).

Next we construct an explicit Euler scheme (3.9) for the fractional diffusion equation with Caputo flux (6.4). For $x_j = jh$, $t_k = k\Delta t$, and $u_j^k = u(x_j, t_k)$, the Grünwald approximation of the Patie–Simon fractional derivative is

$$\mathbb{D}_{[0,x]}^\alpha u_j^k \approx h^{-\alpha} \left[\sum_{i=0}^{j+1} g_{j-i+1}^\alpha u_i^k - g_{j+1}^{\alpha-1} u_0^k \right] = h^{-\alpha} \sum_{i=0}^n b_{ij} u_i^k$$

where $b_{0j} = g_{j+1}^\alpha - g_{j+1}^{\alpha-1}$, $b_{ij} = g_{j-i+1}^\alpha$ for $0 < i \leq j+1$, and $b_{ij} = 0$ for $i > j+1$. Hence the only change in the iteration matrix $B = [b_{ij}]$ is in the top row. From (4.1) it follows easily that

$$g_n^\alpha - g_n^{\alpha-1} = -g_{n-1}^{\alpha-1}, \tag{6.9}$$

and hence we can write $b_{0j} = -g_j^{\alpha-1}$. Now in order to solve the fractional diffusion equation with Caputo flux (6.4), we need only to enforce appropriate boundary conditions.

First assume zero boundary conditions. As in (3.10) it is sufficient to set $b_{ij} = 0$ for $j = 0$ or $j = n$, since a mass proportional to b_{ij} is transported from location x_i to location x_j , and we want this mass to vanish. Then, we obtain the explicit Euler scheme (3.9) with weights

$$b_{ij} = \begin{cases} g_{j-i+1}^\alpha & \text{if } 0 < j < n \text{ and } 0 < i \leq j+1, \\ -g_j^{\alpha-1} & \text{if } i = 0 \text{ and } 0 < j < n, \\ 0 & \text{otherwise.} \end{cases} \tag{6.10}$$

The iteration matrix $B = [b_{ij}]$ differs from (3.10) only in the first row $i = 0$. Since the mass at the left endpoint $x_0 = 0$ is always zero in this case, and since the first row of the matrix B transfers mass out of state $x_0 = 0$, there is no difference in the solutions, and hence Fig. 1 is also the solution to the fractional diffusion equation with Caputo flux (6.4) and absorbing boundary conditions (3.1). In fact, since we assume a zero boundary condition on the left, $u(0, t) = 0$ for all $t > 0$, the fractional diffusion equation with Caputo flux (6.4) and the Riemann–Liouville equation (2.3) on $0 \leq x \leq 1$ are equivalent, due to the relation (6.6).

Next consider a reflecting boundary condition on both sides. Since the iteration matrix $B = [b_{ij}]$ has not changed except in the first row, the argument in Section 4 applies for every state x_j with $j > 0$, i.e., we set $b_{jn} = -g_{n-i}^{\alpha-1}$ for all $i = 1, 2, \dots, n$ as in (4.2). As for the first row, the only way that mass can move to the left boundary $x_0 = 0$ in this scheme is from the adjacent node $x_1 = h$, hence we leave $b_{10} = g_0^\alpha = 1$. In the scheme (3.9) for the Patie–Simon fractional derivative, we have $b_{00} = -g_0^{\alpha-1} = -1$. To prevent mass from state $x_0 = 0$ jumping through the right boundary $x_n = 1$, since $b_{0j} = -g_j^{\alpha-1}$ for $j = 1, 2, \dots, n-1$, and since we require $\sum_j b_{ij} = 0$ for a mass-preserving scheme, we must set

$$b_{0n} = -\sum_{j=0}^{n-1} b_{0j} = 1 + \sum_{j=1}^{n-1} g_j^{\alpha-1} = \sum_{j=0}^{n-1} g_j^{\alpha-1} = g_{n-1}^{\alpha-2}$$

using (4.1). Hence the explicit Euler scheme for this problem is (3.9) with

$$b_{ij} = \begin{cases} g_{j-i+1}^\alpha & \text{if } 0 < j < n \text{ and } 0 < i \leq j+1, \\ 1 & \text{if } i = 1 \text{ and } j = 0, \\ -1 & \text{if } i = j = 0, \\ -g_j^{\alpha-1} & \text{if } i = 0 \text{ and } 0 < j < n, \\ g_{n-1}^{\alpha-2} & \text{if } j = n \text{ and } i = 0, \\ -g_{n-i}^{\alpha-1} & \text{if } j = n \text{ and } 0 < i \leq n, \\ 0 & \text{otherwise.} \end{cases} \tag{6.11}$$




Fig. 5. Numerical solution of the fractional diffusion equation with Caputo flux (6.4) with $\alpha = 1.5$ and $C = 1$ on $0 \leq x \leq 1$ with reflecting boundary conditions (6.12) at time $t = 0$ (solid line), $t = 0.05$ (dashed), $t = 0.1$ (dash dot), $t = 0.5$ (dotted).

Next we will argue that the reflecting boundary conditions for the fractional diffusion equation with Caputo flux (6.4) on $0 \leq x \leq 1$ can be written in the form

$$\partial_{[0,x]}^{\alpha-1}u(0, t) = \partial_{[0,x]}^{\alpha-1}u(1, t) = 0 \quad \text{for all } t \geq 0, \tag{6.12}$$

using the Caputo derivative (6.1). That is, the reflecting boundary conditions zero out the Caputo flux at the boundary. First consider the right boundary $x_n = 1$, and write out the iteration equation for this node: From (3.9) and (6.11) with $\beta = Ch^{-\alpha}\Delta t$ we have

$$u_n^{k+1} = u_n^k - \beta g_{n-1}^{\alpha-2}u_0^k - \beta g_{n-1}^{\alpha-1}u_1^k - \dots - \beta g_1^{\alpha-1}u_{n-1}^k - \beta g_0^{\alpha-1}u_n^k.$$

Using (6.9) this is equivalent to

$$h \frac{u_n^{k+1} - u_n^k}{\Delta t} = -Ch^{1-\alpha} \left[\sum_{i=0}^n g_{n-i}^{\alpha-1}u_i^k - g_n^{\alpha-2}u_0^k \right].$$

Letting $\Delta t \rightarrow 0$ and $h \rightarrow 0$, the left-hand side converges to zero, the first term on the right converges to $\mathbb{D}_{[0,x]}^{\alpha-1}u(1, t)$ using the Grünwald approximation (3.4), and recalling that $hn = 1$, the second term

$$Ch^{1-\alpha}g_n^{\alpha-2}u_0^k \sim Cn^{\alpha-1} \frac{2-\alpha}{\Gamma(3-\alpha)}n^{1-\alpha}u_0^k \rightarrow \frac{C}{\Gamma(2-\alpha)}u(0, t)$$

as $h \rightarrow 0$ using (6.8). Using (6.2) with $\gamma = \alpha - 1$ and $x = 1$, it follows that the entire right-hand side converges to the Caputo derivative of order $\alpha - 1$, and hence the reflecting boundary condition (6.12) holds at the right boundary $x = 1$.

Using $b_{0j} = g_{j+1}^\alpha - g_{j+1}^{\alpha-1}$, the iteration equation at the left boundary is $u_0^{k+1} = u_0^k + \beta g_1^\alpha u_0^k - \beta g_1^{\alpha-1} u_0^k + \beta g_0^{\alpha-1} u_1^k$. This reduces to

$$h \frac{u_0^{k+1} - u_0^k}{\Delta t} = -Ch^{1-\alpha} \left[\sum_{i=0}^1 g_{1-i}^{\alpha-1}u_i^k - g_1^{\alpha-2}u_0^k \right]$$

which is consistent with the reflecting boundary condition (6.12) at the left boundary $x = 0$. A rigorous proof that the left boundary condition in (6.12) holds is similar to the case of the Riemann–Liouville generator, see [76, Proposition 19].

Remark 6.1. Comparing (4.3) and (6.12) shows that the form of the reflecting boundary condition *also changes* when we change the type of fractional derivative in the fractional diffusion equation. When $\alpha = 2$, both forms reduce to the classical reflecting boundary condition $\frac{\partial}{\partial x}u(0, t) = \frac{\partial}{\partial x}u(1, t) = 0$. D. del-Castillo-Negrete [56, Section II.B] suggested using a Caputo fractional flux to avoid singularities at the boundary.

Fig. 5 shows a numerical solution of the fractional diffusion equation with Caputo flux (6.4) on $0 \leq x \leq 1$ with reflecting boundary conditions, using the same numerical method and initial function as in Fig. 1. Solution curves are skewed for $0 < t < \infty$, and the total mass remains equal to the initial mass $M = 1$ for all $t > 0$, since the scheme is mass-preserving. As t increases, the solutions approach the unique steady state solution $u_\infty(x) = 1$ on $0 \leq x \leq 1$ with unit mass, which is




Fig. 6. Numerical solution of the fractional diffusion equation with Caputo flux (6.4) with $\alpha = 1.5$ and $C = 1$ on $0 \leq x \leq 1$ with boundary conditions (6.14): reflecting on the left, and absorbing on the right at time $t = 0$ (solid line), $t = 0.05$ (dashed), $t = 0.1$ (dash dot), $t = 0.5$ (dotted).

very different than the unit mass steady state solution $u_\infty(x) = (\alpha - 1)x^{\alpha-2}$ to the Riemann–Liouville fractional diffusion equation (2.3) on $0 \leq x \leq 1$ with reflecting boundary conditions. In the present case, the steady state solution is easy to verify, by simply plugging into (6.4).

Remark 6.2. The left reflecting boundary condition for the fractional diffusion equation with Caputo flux (6.4) can also be written in the traditional form

$$\frac{\partial}{\partial x} u(0, t) = 0 \quad \text{for all } t \geq 0. \tag{6.13}$$

To see this, rewrite the iteration equation $u_0^{k+1} = u_0^k - \beta u_0^k + \beta u_1^k$ in the equivalent form

$$h^{\alpha-1} \frac{u_n^{k+1} - u_n^k}{\Delta t} = C \left[\frac{u_1^k - u_0^k}{h} \right]$$

and let $h \rightarrow 0$ and $\Delta t \rightarrow 0$. In view of (6.1), the condition (6.13) also implies that the Caputo fractional derivative $\partial_{[0,x]}^{\alpha-1} u(0, t) = 0$. The same is *not* true of the Riemann–Liouville derivative, and indeed, even the steady state solution $u_\infty(x) = (\alpha - 1)x^{\alpha-2}$ of the Riemann–Liouville fractional diffusion equation (2.3) with reflecting boundary conditions does not satisfy the condition (6.13). The nonlocal right reflecting boundary condition for the fractional diffusion equation with Caputo flux (6.4) cannot be reduced to a local first derivative condition, since it depends on the values of the solution across the entire domain. Indeed, one can see in Fig. 5 that $\frac{\partial}{\partial x} u(1, t) \neq 0$ at the right boundary.

Next we consider the fractional diffusion equation with Caputo flux (6.4) on $0 \leq x \leq 1$ with boundary conditions

$$\partial_{[0,x]}^{\alpha-1} u(0, t) = 0 \quad \text{and} \quad u(1, t) = 0 \quad \text{for all } t \geq 0, \tag{6.14}$$

reflecting at the left boundary $x = 0$ and absorbing at the right boundary $x = 1$. Here we simply zero out the coefficients b_{ij} from (6.11) governing mass transport from state i to state $j = n$. This yields the explicit Euler scheme (3.9) with

$$b_{ij} = \begin{cases} g_{j-i+1}^\alpha & \text{if } 0 < j < n \text{ and } 0 < i \leq j + 1, \\ 1 & \text{if } i = 1 \text{ and } j = 0, \\ -1 & \text{if } i = j = 0, \\ -g_j^{\alpha-1} & \text{if } i = 0 \text{ and } 0 < j < n, \\ 0 & \text{otherwise.} \end{cases} \tag{6.15}$$

Fig. 6 shows a numerical solution of the fractional diffusion equation with Caputo flux (6.4) on $0 \leq x \leq 1$ with boundary conditions (6.14), using the same numerical method and initial function as in Fig. 1. Solution curves are skewed for all $t > 0$, and approach the unique steady state solution $u_\infty(x) = 0$ on $0 \leq x \leq 1$ as t increases. In [76, Proposition 19] it is shown that the Cauchy problem (6.4) with these boundary conditions (6.12) or (6.14) is well-posed on the Banach space $L^1[0, 1]$, and the exact domain of the generator is computed.

Finally we consider the fractional diffusion equation with Caputo flux (6.4) on $0 \leq x \leq 1$ with boundary conditions

$$u(0, t) = 0 \quad \text{and} \quad \partial_{[0,x]}^{\alpha-1} u(1, t) = 0 \quad \text{for all } t \geq 0, \tag{6.16}$$

absorbing at the left boundary $x = 0$ and reflecting at the right boundary $x = 1$. Since $u(0, t) = 0$ for all $t > 0$, this problem is mathematically equivalent to the fractional diffusion equation (2.3) on $0 \leq x \leq 1$ with boundary conditions (5.3). Hence Fig. 4 also represents the solution to this fractional boundary value problem. One can also see this by setting $b_{i0} = 0$ in (6.11) to get

$$b_{ij} = \begin{cases} g_{j-i+1}^\alpha & \text{if } 0 < j < n \text{ and } 0 < i \leq j + 1, \\ -g_j^{\alpha-1} & \text{if } i = 0 \text{ and } 0 < j < n, \\ -g_{n-1}^{\alpha-2} & \text{if } j = n \text{ and } i = 0, \\ -g_{n-i}^{\alpha-1} & \text{if } j = n \text{ and } 0 < i \leq n, \\ 0 & \text{otherwise.} \end{cases} \tag{6.17}$$

Since $u_0^k = 0$ for all k , the first row of the iteration matrix B is immaterial, and the rest of the matrix is exactly the same as for the corresponding case of the fractional diffusion equation (2.3).

Remark 6.3. The general steady state solution to the fractional diffusion equation with Caputo flux (6.4) is $u_\infty(x) = c_1 x^{\alpha-1} + c_2$. This can be verified by a calculation similar to Remark 4.2: Since $\mathbf{D}_{[0,x]}^\alpha u(x) = \frac{d}{dx} \mathbb{J}_{[0,x]}^{2-\alpha} u'(x)$, we have

$$\mathbf{D}_{[0,x]}^\alpha u_\infty(x) = \frac{d}{dx} \mathbb{J}_{[0,x]}^{2-\alpha} c_1(\alpha - 1)x^{\alpha-2} = \frac{d}{dx} [c_1(\alpha - 1)\Gamma(\alpha - 1)] = 0$$

using (4.7). Zero boundary conditions require $c_2 = 0$ to make $u_\infty(0) = 0$, and then also $c_1 = 0$ to make $u_\infty(1) = 0$. For reflecting boundary conditions, we compute

$$\partial_{[0,x]}^{\alpha-1} u_\infty(x) = \mathbb{J}_{[0,x]}^{2-\alpha} u'_\infty(x) = c_1 \Gamma(\alpha)$$

for $0 < x < 1$. The right boundary condition $\partial_{[0,x]}^{\alpha-1} u_\infty(1) = 0$ requires $c_1 = 0$, and then the left boundary condition is satisfied for any real number c_2 . Take $c_2 = 1$ to get the solution with total mass 1. If just the left boundary condition is absorbing, we require $c_2 = 0$, and then the reflecting boundary condition on the right requires $c_1 = 0$ as well. If just the right boundary condition is absorbing, then $c_1 + c_2 = 0$. Then if the left boundary is reflecting, $c_1 = 0$, and hence $c_2 = 0$ as well.

Remark 6.4. Cushman and Ginn [27] use the fractional derivative (6.4) (on the real line, with the lower integration limit 0 changed to $-\infty$) to model contaminant transport in groundwater, see also [80]. For such problems, all three fractional derivatives are equivalent, since the boundary term at $x = -\infty$ vanishes.

Remark 6.5. In [81] we show that the backward generator of a standard spectrally negative α -stable process reflected to stay positive is the Caputo fractional derivative (6.1). Since [79, Eq. (1.2)]

$$\partial_{[0,x]}^\alpha u(x) = \mathbf{D}_{[0,x]}^\alpha u(x) - u'(0) \frac{x^{1-\alpha}}{\Gamma(2-\alpha)} \tag{6.18}$$

for $1 < \alpha < 2$, and since $u'(0) = 0$ for every function in the domain of the generator [79, Remark 2.3 (a)], these two forms are completely equivalent.

Remark 6.6. For either the Riemann–Liouville fractional diffusion equation (2.1) or the fractional diffusion equation with Caputo flux (6.4), the fractional derivative operator with a zero boundary condition at one or both boundaries is invertible [76, Proposition 6]. This implies that, for any initial data $u_0(x)$, the solution converges to the unique steady state solution $u_\infty = 0$, see Appendix for details. In the case of reflecting boundary conditions, one can write the solution $u = (u - u_\infty) + u_\infty$, and apply a similar argument to the first term, to show that all solutions converge to steady state.

7. What can go wrong

One could also consider the Caputo fractional differential equation

$$\frac{\partial}{\partial t} u(x, t) = C \partial_{[0,x]}^\alpha u(x, t) \tag{7.1}$$

with $1 < \alpha < 2$ on the unit interval $0 \leq x \leq 1$, using the Caputo fractional derivative (6.1). However, solutions to (7.1) are not positivity preserving. An explicit Euler scheme to solve this problem can be developed using the Caputo Grünwald formula

$$\begin{aligned} \partial_{[0,x]}^\alpha f(x) = \lim_{h \rightarrow 0} h^{-\alpha} & \left[\sum_{i=0}^{j+1} g_i^\alpha f(x - (i-1)h) - g_{j+1}^{\alpha-1} f(x - jh) \right. \\ & \left. - g_{j+1}^{\alpha-2} f(x - (j-1)h) + g_{j+1}^{\alpha-2} f(x - jh) \right] \end{aligned} \tag{7.2}$$




Fig. 7. Numerical solution of the Caputo fractional differential equation (7.1) with $\alpha = 1.5$ and $C = 1$ on $0 \leq x \leq 1$ with zero boundary conditions (3.1) at time $t = 0$ (solid line), $t = 0.01$ (dashed), $t = 0.04$ (dash dot), $t = 0.2$ (dotted). Solutions take negative values even though the initial function (7.4) is nonnegative.

where $j = j(h) = \lceil x/h \rceil$. The proof that (7.2) holds is very similar to (6.7). This leads to the explicit Euler scheme (3.9) with

$$b_{ij} = \begin{cases} g_{j-i+1}^\alpha & \text{if } 0 < j < n \text{ and } 1 < i \leq j + 1, \\ -g_j^{\alpha-1} + g_{j+1}^{\alpha-2} & \text{if } i = 0 \text{ and } 0 < j < n, \\ g_j^\alpha - g_{j+1}^{\alpha-2} & \text{if } i = 1 \text{ and } 0 < j < n, \\ 0 & \text{otherwise.} \end{cases} \tag{7.3}$$

Fig. 7 shows a numerical solution of the fractional differential equation (7.1) on $0 \leq x \leq 1$ with Dirichlet boundary conditions (3.1), and initial function

$$u_0(x) = \begin{cases} \frac{64\pi^3}{\pi^2 - 4} \left(x - \frac{1}{4}\right)^2 \sin(4\pi x) & \text{for } 0 < x < 0.25, \\ 0 & \text{otherwise,} \end{cases} \tag{7.4}$$

using the same numerical method as in Fig. 1. Since the solution takes negative values with nonnegative initial data, the Caputo fractional differential equation (7.1) cannot provide a physically meaningful model for anomalous diffusion.

Remark 7.1. If one considers the Caputo fractional differential equation (7.1) on $0 \leq x \leq 1$ with $1 < \alpha < 2$ and the traditional Neumann boundary conditions

$$\frac{\partial}{\partial x} u(0, t) = \frac{\partial}{\partial x} u(1, t) = 0 \quad \text{for all } t \geq 0, \tag{7.5}$$

then the Caputo and Patie–Simon fractional derivatives are equal, in light of (6.18). Since (7.5) implies (6.12) by (6.1), solutions to (7.1) with the boundary conditions (7.5) also solve the problem (6.4) with reflecting boundary conditions (6.12). However, the domain of the fractional derivative (6.5) with the reflecting boundary conditions (6.12) is strictly larger, and there are solutions to (6.4) with reflecting boundary conditions (6.12) that do not solve (7.1) with the boundary conditions (7.5), e.g., note that $\frac{\partial}{\partial x} u(1, t) \neq 0$ in Fig. 5.

Acknowledgments

The authors would like to thank James F. Kelly, Department of Statistics and Probability, Michigan State University, for helpful discussions.

Baeumer and Kovács were partially supported by the Marsden Fund administered by the Royal Society of New Zealand (UO01418). Meerschaert was partially supported by ARO MURI grant W911NF-15-1-0562 and NSF (National Science Foundation (USA)) grants DMS-1462156 and EAR-1344280. Sankaranarayanan was supported by ARO MURI grant W911NF-15-1-0562.

Appendix

The following result implies that solutions in this paper converge to the steady state solution $u_\infty = 0$ if at least one boundary condition is absorbing. This follows because, in this case, the fractional derivative operator is invertible [76, Proposition 6] and generates a strongly continuous positive contraction semigroup [76, Theorem 20].

Lemma A.1. *Let (Ω, μ) be a σ -finite measure space and $X = L^p(\Omega)$, $1 \leq p < \infty$, or let Ω be a locally compact Hausdorff space and $X = C_0(\Omega)$. Suppose that A generates a strongly continuous positive contraction semigroup on X and that A^{-1} exists as a bounded operator on X . Then, for all $x \in X$, we have $\|T(t)x\| \rightarrow 0$ as $t \rightarrow \infty$ exponentially fast.*

Proof. Let $\sigma(A)$ denote the spectrum of A and $\rho(A)$ the resolvent set of A . Since A generates a strongly continuous contraction semigroup, it follows from the Hille–Yosida Theorem that $(0, \infty) \subset \rho(A)$. As A^{-1} exists as a bounded operator on X , it follows that $0 \in \rho(A)$. Since A generates a strongly continuous positive contraction semigroup on X , it follows that the resolvent $R(\lambda, A)$ of A satisfies $R(\lambda, A) = \int_0^\infty e^{-\lambda t} T(t) dt \geq 0$ (strong Bochner integral) for $\lambda > 0$ as the positive cone is closed in X . The resolvent is an analytic function of λ for $\lambda \in \rho(A)$ and hence continuous. Therefore $A^{-1} = \lim_{\lambda \rightarrow 0^+} R(\lambda, A) \geq 0$, again, since the positive cone is closed in X .

Let $s(A)$ denote the spectral bound of A ; that is,

$$s(A) = \sup\{\operatorname{Re} \lambda : \lambda \in \sigma(A)\}$$

and $\omega_0 \in \mathbb{R}$ the growth bound of the semigroup; that is,

$$\omega_0 = \inf\{\omega \in \mathbb{R} : \text{there is } M_\omega \geq 1 \text{ such that } \|T(t)\| \leq M_\omega e^{\omega t}, \quad t \geq 0\}.$$

It follows from [82, Chapter VI, Lemma 1.9] that $A^{-1} \geq 0$ implies that $s(A) < 0$. Finally, by [83, Theorem 5.3.6] when $X = L^p(\Omega)$ and [83, Theorem 5.3.8] when $X = C_0(\Omega)$, it follows that $\omega_0 = s(A)$ and hence $\omega_0 < 0$. Thus, there is $\epsilon > 0$ and $M_\epsilon \geq 1$ such that $\|T(t)\| \leq M_\epsilon e^{-\epsilon t}$, $t \geq 0$, and the proof is complete. \square

References

- [1] R. Herrmann, *Fractional Calculus: An Introduction for Physicists*, second ed., World Scientific, Singapore, 2014.
- [2] R. Klages, G. Radons, I.M. Sokolov, *Anomalous Transport: Foundations and Applications*, Wiley-VCH, Weinheim, Germany, 2008.
- [3] F. Mainardi, *Fractional Calculus and Waves in Linear Viscoelasticity: An Introduction to Mathematical Models*, World Scientific, 2010.
- [4] F. Mainardi, *Fractals and Fractional Calculus in Continuum Mechanics*, Springer, Berlin, 1997.
- [5] R. Metzler, J. Klafter, The random walk's guide to anomalous diffusion: a fractional dynamics approach, *Phys. Rep.* 339 (1) (2000) 1–77.
- [6] R. Metzler, J. Klafter, The restaurant at the end of the random walk: recent developments in the description of anomalous transport by fractional dynamics, *J. Phys. A* 37 (31) (2004) R161–R208.
- [7] I. Podlubny, *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of their Solution and Some of their Applications*, Academic Press, San Diego, California, 1999.
- [8] S. Samko, A. Kilbas, O. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*, Gordon and Breach, London, 1993.
- [9] R. Hilfer, *Applications of Fractional Calculus in Physics*, World Scientific, 2000.
- [10] M. Magdziarz, A. Weron, K. Weron, Fractional Fokker–Planck dynamics: Stochastic representation and computer simulation, *Phys. Rev. E* 75 (2007) 016708.
- [11] A. Piryatinska, A.I. Saichev, W.A. Woyczynski, Models of anomalous diffusion: Subdiffusive and superdiffusive cases, *Physica A* 349 (2005) 375–424.
- [12] V.E. Tarasov, Fractional vector calculus and fractional Maxwell's equations, *Ann. Phys.* 323 (2008) 2756–2778.
- [13] R. Gorenflo, F. Mainardi, E. Scalas, M. Raberto, Fractional Calculus and Continuous-Time Finance. III. The Diffusion Limit. *Mathematical Finance* (Konstan, 2000), in: *Trends Math.*, Birkhäuser, Basel, 2001, pp. 171–180.
- [14] A. Jurlewicz, A. Wyłomańska, P. Żebrowski, Coupled continuous-time random walk approach to the Rachev–Rüschendorf model for financial data, *Physica A* 388 (2009) 407–418.
- [15] F. Mainardi, M. Raberto, R. Gorenflo, E. Scalas, Fractional Calculus and continuous-time finance II: the waiting-time distribution, *Physica A* 287 (2000) 468–481.
- [16] E. Scalas, R. Gorenflo, F. Mainardi, Fractional calculus and continuous-time finance, *Physica A* 284 (2000) 376–384.
- [17] E. Scalas, Five years of continuous-time random walks in econophysics, in: *The Complex Networks of Economic Interactions*, Springer, Berlin, 2006, pp. 3–16.
- [18] E. Barkai, Y. Garini, R. Metzler, Strange kinetics of single molecules in living cells, *Phys. Today* 65 (2012) 29.
- [19] S. Fedotov, A. Iomin, Migration and proliferation dichotomy in tumor-cell invasion, *Phys. Rev. Lett.* 98 (2007) 8101.
- [20] J.-H. Jeon, H. Martinez-Seara Monne, M. Javanainen, R. Metzler, Anomalous diffusion of phospholipids and cholesterol in a lipid bilayer and its origins, *Phys. Rev. Lett.* 109 (2012) 188103.
- [21] W. Chen, S. Holm, Fractional Laplacian time-space models for linear and nonlinear lossy media exhibiting arbitrary frequency power-law dependency, *J. Acoust. Soc. Am.* 115 (2004) 1424–1430.
- [22] R.L. Magin, *Fractional Calculus in Bioengineering*, Begell House, 2006.
- [23] R.L. Magin, C. Ingo, L. Colon-Perez, W. Triplett, T.H. Mareci, Characterization of anomalous diffusion in porous biological tissues using fractional order derivatives and entropy, *Microporous Mesoporous Mater.* 71 (2014) 617–627.
- [24] M. Palombo, A. Gabrielli, S. De Santis, C. Cametti, G. Ruocco, S. Capuani, Spatio-temporal anomalous diffusion in heterogeneous media by nuclear magnetic resonance, *J. Chem. Phys.* 135 (3) (2011) 034504.
- [25] B.E. Treeby, B.T. Cox, Modeling power law absorption and dispersion for acoustic propagation using the fractional Laplacian, *J. Acoust. Soc. Am.* 127 (2010) 2741–2748.
- [26] X.J. Zhou, Q. Gao, O. Abdullah, R.L. Magin, Studies of anomalous diffusion in the human brain using fractional order calculus, *Magn. Reson. Med.* 63 (2010) 562–569.

- [27] J.H. Cushman, T. Ginn, Fractional advection–dispersion equation: A classical mass balance with convolution–Fickian flux, *Water Resour. Res.* 36 (2000) 3763–3766.
- [28] Z.-Q. Deng, L. Bengtsson, V.P. Singh, Parameter estimation for fractional dispersion model for rivers, *Environ. Fluid Mech.* 6 (2006) 451–475.
- [29] S. Kim, M.L. Kavvas, Generalized Fick's law and fractional ADE for pollutant transport in a river: detailed derivation, *J. Hydrol. Eng.* 11 (2006) 80–83.
- [30] C. Shen, M.S. Phanikumar, An efficient space-fractional dispersion approximation for stream solute transport modeling, *Adv. Water Resour.* 32 (2009) 1482–1494.
- [31] G. Zaslavsky, Fractional kinetic equation for Hamiltonian chaos, *Physica D* 76 (1994) 110–122.
- [32] G. Zaslavsky, Chaos, fractional kinetics, and anomalous transport, *Phys. Rep.* 371 (2002) 461–580.
- [33] M. Gosak, M. Marhl, M. Perc, Chaos out of internal noise in the collective dynamics of diffusively coupled cells, *Eur. Phys. J. B* 62 (2008) 171–177.
- [34] M. Gosak, A. Stožer, R. Markovič, J. Dolenšek, M. Marhl, M. Slak Rupnik, M. Perc, The relationship between node degree and dissipation rate in networks of diffusively coupled oscillators and its significance for pancreatic beta cells, *Chaos* 25 (2015) 073115.
- [35] X. Sun, M. Perc, Q. Lu, J. Kurths, Spatial coherence resonance on diffusive and small-world networks of Hodgkin–Huxley neurons, *Chaos* 18 (2008) 023102.
- [36] A.H. Bhrawy, M.A. Zaky, A method based on the Jacobi tau approximation for solving multi-term time-space fractional partial differential equations, *J. Comput. Phys.* 281 (2015) 876–895.
- [37] Z. Deng, V.P. Singh, L. Bengtsson, Numerical solution of fractional advection–dispersion equation, *J. Hydraul. Eng.* 130 (2004) 422–431.
- [38] K. Diethelm, N.J. Ford, A.D. Freed, A predictor–corrector approach for the numerical solution of fractional differential equations, *Nonlinear Dynam.* 29 (1–4) (2002) 3–22.
- [39] G.J. Fix, J.P. Roop, Least squares finite element solution of a fractional order two-point boundary value problem, *Comput. Math. Appl.* 48 (2004) 1017–1033.
- [40] F. Liu, V. Ahn, I. Turner, Numerical solution of the space fractional Fokker–Planck equation, *J. Comput. Appl. Math.* 166 (2004) 209–219.
- [41] V.E. Lynch, B.A. Carreras, D. del Castillo–Negrete, K.M. Ferreira–Mejias, H.R. Hicks, Numerical methods for the solution of partial differential equations of fractional order, *J. Comput. Phys.* 192 (2003) 406–421.
- [42] M.M. Meerschaert, C. Tadjeran, Finite difference approximations for fractional advection–dispersion flow equations, *J. Comput. Appl. Math.* 172 (2004) 65–77.
- [43] M.M. Meerschaert, C. Tadjeran, Finite difference approximations for two-sided space-fractional partial differential equations, *Appl. Numer. Math.* 56 (2006) 80–90.
- [44] M.M. Meerschaert, H.P. Scheffler, C. Tadjeran, Finite difference methods for two-dimensional fractional dispersion equation, *J. Comput. Phys.* 211 (2006) 249–261.
- [45] Z. Odibat, S. Momani, Numerical methods for nonlinear partial differential equations of fractional order, *Appl. Math. Model.* 32 (1) (2008) 28–39.
- [46] I. Podlubny, A. Chechkin, T. Skovranek, Y. Chen, B.M.V. Jara, Matrix approach to discrete fractional calculus ii: Partial fractional differential equations, *J. Comput. Phys.* 228 (2009) 3137–3153.
- [47] C. Tadjeran, M.M. Meerschaert, H.P. Scheffler, A second order accurate numerical approximation for the fractional diffusion equation, *J. Comput. Phys.* 213 (2006) 205–213.
- [48] H. Wang, D. Yang, S. Zhu, Inhomogeneous Dirichlet boundary-value problems of space-fractional diffusion equations and their finite element approximations, *SIAM J. Numer. Anal.* 52 (2014) 1292–1310.
- [49] S.B. Yuste, L. Acedo, An explicit finite difference method and a new von Neumann type stability analysis for fractional diffusion equations, *SIAM J. Numer. Anal.* 42 (2005) 1862–1874.
- [50] H. Zhang, F. Liu, M.S. Phanikumar, M.M. Meerschaert, A novel numerical method for the time variable fractional order mobile–immobile advection–dispersion model, *Comput. Math. Appl.* 66 (5) (2013) 693–701.
- [51] B. Baeumer, T. Luks, M.M. Meerschaert, Space–time fractional Dirichlet problems. Preprint at www.stt.msu.edu/users/mcubed/SpaceTimeFrac.pdf.
- [52] O. Defterli, M. D'Elia, Q. Du, M. Gunzburger, R. Lehoucq, M.M. Meerschaert, Fractional diffusion on bounded domains, *Fract. Calc. Appl. Anal.* 18 (2) (2015) 342–360.
- [53] B. Jin, R. Lazarov, J. Pasciak, W. Rundell, Variational formulation of problems involving fractional order differential operators, *Math. Comp.* 84 (2015) 2665–2700.
- [54] R.A.C. Ferreira, A Lyapunov-type inequality for a fractional boundary value problem, *Fract. Calc. Appl. Anal.* 16 (2013) 978–984.
- [55] X. Zhu, Y. Feng, Y. Wang, A Lyapunov-type inequality for a periodic boundary value problem of a fractional differential equation, *Ann. Appl. Math.* 33 (2017) 212–220.
- [56] D. del Castillo–Negrete, Fractional diffusion models of nonlocal transport, *Phys. Plasmas* 13 (2006) 082308.
- [57] E. Montefusco, B. Pellacci, G. Verzini, Fractional diffusion with Neumann boundary conditions: the logistic equation, *Discrete Contin. Dyn. Syst. Ser. B* 18 (2013) 2175–2202.
- [58] J. Xie, Q. Huang, F. Zhao, H. Gui, Block pulse functions for solving fractional Poisson type equations with Dirichlet and Neumann boundary conditions, *Bound. Value Probl.* 2017 (2017) 32.
- [59] N. Laskin, Fractals and quantum mechanics, *Chaos* 10 (4) (2000) 780–790.
- [60] B. Szekeres, F. Izsák, A finite difference method for fractional diffusion equations with Neumann boundary conditions, *Open Math.* 13 (2015) 581–600.
- [61] S.C. Lim, L.P. Teo, Finite temperature Casimir effect in piston geometry and its classical limit, *Eur. Phys. J. C* 60 (2009) 323–344.
- [62] S.C. Lim, L.P. Teo, Repulsive Casimir force from fractional Neumann boundary conditions, *Phys. Lett. B* 679 (2009) 130–137.
- [63] H. Wang, D. Yang, Wellposedness of Neumann boundary-value problems of space-fractional differential equations, Preprint arxiv.org/abs/1612.02450v1.
- [64] V.R. Voller, An exact solution of a limit case Stefan problem governed by a fractional diffusion equation, *Int. J. Heat Mass Transfer* 53 (2010) 5622–5625.
- [65] V. Kolokoltsov, On fully mixed and multidimensional extensions of the Caputo and Riemann–Liouville derivatives, related Markov processes and fractional differential equations, *Fract. Calc. Appl. Anal.* 18 (2015) 1039–1073.
- [66] M. Warma, The fractional relative capacity and the fractional Laplacian with Neumann and Robin boundary conditions on open sets, *Potential Anal.* 42 (2015) 499–547.
- [67] S. Umarov, Fractional Fokker–Planck–Kolmogorov equations associated with SDES on a bounded domain, *Fract. Calc. Appl. Anal.* 20 (2017) 1281–1304.
- [68] Q. Du, M. Gunzburger, R.B. Lehoucq, K. Zhou, Analysis and approximation of nonlocal diffusion problems with volume constraints, *SIAM Rev.* 54 (2012) 667–696.
- [69] N. Krepyshcheva, L. Di Pietro, M.C. Néel, Space-fractional advection–diffusion and reflective boundary condition, *Phys. Rev. E* 73 (2006) 021104.
- [70] N. Cusimano, K. Burrage, I. Turner, D. Kay, On reflecting boundary conditions for space-fractional equations on a finite interval: Proof of the matrix transfer technique, *Appl. Math. Model.* 42 (2017) 554–565.
- [71] M. Ilic, F. Liu, I. Turner, V. Anh, Numerical approximation of a fractional-in-space diffusion equation, I, *Fract. Calc. Appl. Anal.* 8 (2005) 323–341.
- [72] M.M. Meerschaert, A. Sikorskii, Stochastic Models for Fractional Calculus, in: De Gruyter Studies in Mathematics, vol. 43, De Gruyter, Berlin, 2012.
- [73] M.M. Meerschaert, Mathematical Modeling, fourth ed., Academic Press, 2013.

- [74] B. Baeumer, M. Kovács, H. Sankaranarayanan, Higher order Grunwald approximations of fractional derivatives and fractional powers of operators, *Trans. Amer. Math. Soc.* 367 (2015) 813–834.
- [75] R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2017. <http://www.R-project.org/>.
- [76] B. Baeumer, M. Kovács, H. Sankaranarayanan, Fractional partial differential equations with boundary conditions, *J. Differential Equations* 264 (2018) 1377–1410.
- [77] R.D. Richtmyer, K.W. Morton, *Difference Methods for Initial-Value Problems*, Krieger Publishing, Malabar, Florida, 1994.
- [78] R. Schumer, D.A. Benson, M.M. Meerschaert, S.W. Wheatcraft, Eulerian derivation for the fractional advection–dispersion equation, *J. Contam. Hydrol.* 48 (2001) 69–88.
- [79] P. Patie, T. Simon, Intertwining certain fractional derivatives, *Potential Anal.* 36 (2012) 569–587.
- [80] M.M. Meerschaert, J. Mortensen, S.W. Wheatcraft, Fractional vector calculus for fractional advection–dispersion, *Physica A* 367 (2006) 181–190.
- [81] B. Baeumer, M. Kovács, M.M. Meerschaert, R.L. Schilling, P. Straka, Reflected spectrally negative stable processes and their governing equations, *Trans. Amer. Math. Soc.* 368 (1) (2016) 227–248.
- [82] K.J. Engel, R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Springer-Verlag, 2000.
- [83] W. Arendt, C.J.K. Batty, M. Hieber, F. Neubrander, *Vector-Valued Laplace Transforms and Cauchy Problems*, second ed., in: *Monographs in Mathematics*, vol. 96, Birkhäuser Verlag, Basel, 2010.