

1-2. A person will be selected at random from all those sorted by sex and gene type in the table below.

	AA	Aa	aa	
male	5	7	5	17
female	6	9	3	18
				35

1. What is the conditional probability  $P(aa \mid \text{if female})$ , that the selected person has gene type aa if they are female?

- a.  $\frac{5}{17}$  b.  $\frac{17}{35}$  c.  $\frac{3}{18}$  d.  $\frac{25}{35}$  e.  $\frac{27}{35}$   $\frac{3}{18}$  6 9 (3) 18

2. What is the probability the person does not have gene type aa?

- a.  $\frac{5}{17}$  b.  $\frac{17}{35}$  c.  $\frac{3}{18}$  d.  $\frac{25}{35}$  e.  $\frac{27}{35}$  57 69 total 27

3. The number  $X$  of insurance claims for broken windshield averages 400 per week. There are many opportunities for such a claim to be made and each has a small probability of happening. These appear to be independent events so the Poisson model is deemed appropriate for random variable  $X$ . Determine a 95% interval for  $X$  (answer rounded).

- a. [380, 420] b. [180, 220] c. [0, 800] d. [360, 440] e. [300, 500]

$\mu = \lambda = 400$   
 $\sigma = \sqrt{\mu} = \sqrt{\lambda} = 20$

4-7. Suppose that  $P(OIL) = 0.4$ ,  $P(- \mid \text{if } OIL) = 0.3$ ,  $P(- \mid \text{if } OIL^c) = 0.8$ .

4.  $P(OIL^c -) = P(OIL^c)P(- \mid OIL^c) = 0.6 \cdot 0.8 = 0.48$

- a. 0.8 b. 0.6 c. 0.48 d. 0.2 e. 0.3

5.  $P(-) = P(OIL-) + P(OIL^c-) = 0.4 \cdot 0.3 + 0.6 \cdot 0.8 = 0.12 + 0.48 = 0.6$

- a. 0.8 b. 0.6 c. 0.48 d. 0.2 e. 0.3

6.  $P(OIL \mid \text{if } -) = \frac{P(OIL-)}{P(-)} = \frac{0.12}{0.6} = 0.2$

- a. 0.8 b. 0.6 c. 0.48 d. 0.2 e. 0.3

Suppose also that the cost to test is 10, the cost to drill is 60, and the gross return from finding OIL is 400.

7. NET return from policy "test, but only drill if the test is positive" for the contingency

$OIL^c +$  is equal to:

- a. 340 b. 330 c. -70 d. -100 e. -60

TEST	DRILL	OIL	RETURN
-10	-60	0 OR OIL <sup>c</sup>	

8. People are equal-probability selected from a population and sorted by sex and gene type. Which one of the tables below exhibits independence between sex and gene type?

a.		AA	Aa	aa
	male	20	15	10
	female	50	45	20
b.		AA	Aa	aa
	male	40	40	20
	female	60	90	30
c.		AA	Aa	aa
	male	60	30	20
	female	90	45	30
d.		AA	Aa	aa
	male	20	10	10
	female	30	45	15
e.		AA	Aa	aa
	male	60	45	15
	female	20	65	5

60:90 or 30:45 or 20:30  
 PROPORTIONALITY IS THE  
 APPEARANCE OF INDEPENDENCE  
 IN TREEED COUNTS.

9-11. We are given the probability distribution and summary totals shown:

x	p(x)	x p(x)	x <sup>2</sup> p(x)
-2	0.1	-0.2	0.4
0	0.7	0	0
4	0.2	0.8	3.2

9.  $E X^2 = \sum x^2 p(x) = 3.6$   
 a. ~7   b. 15   c. 2.8   d. 3.6   e. 3.24

totals 1.0

0.6  
EX  
3.6  
E(X<sup>2</sup>)

10. Var X = (you may use a calculator or employ summary information above)

a. ~7   b. 15   c. 2.8   d. 3.6   e. 3.24

Var X =  $E(X^2) - (EX)^2 = 3.6 - 0.6^2$   
 SHORT FORM = 3.24

11. P(X is larger or equal 0) =

a. 0.5   b. 1   c. 0.2   d. 0.9   e. 0.3

ONLY 0 AND 4 ARE  $\geq 0$

$P(0) = 0.7$   
 $P(4) = 0.2$  / and 0.9

12-14. A plane has two engines. The probability of the left engine failing is 0.002. The probability of the right engine failing is 0.003. Engine failures are independent events.

12. What is the probability that (the left engine fails)  $\cap$  and (the right does not)?

- a.  $0.002 + 0.003 - 0.002 \cdot 0.003$  b.  $0.002 \cdot 0.003$  c.  $0.998 \cdot 0.003$  d.  $0.002 \cdot 0.997$  e.  $0.998 \cdot 0.997$

$L \text{ FAILS } R \text{ NOT } = 0.002 \cdot 0.997$

13. What is the probability that (the left engine fails)  $\cup$  or (the right engine fails)?

- a.  $0.002 + 0.003 - 0.002 \cdot 0.003$  b.  $0.002 \cdot 0.003$  c.  $0.998 \cdot 0.003$  d.  $0.002 \cdot 0.997$  e.  $0.998 \cdot 0.997$

ADDITION RULE

14. What is the probability that neither engine fails?

- a.  $0.002 + 0.003 - 0.002 \cdot 0.003$  b.  $0.002 \cdot 0.003$  c.  $0.998 \cdot 0.003$  d.  $0.002 \cdot 0.997$  e.  $0.998 \cdot 0.997$

15-19. A lottery has return  $X$  which is a random variable with

$E X = 2$        $\text{Var } X = 9$

15. Variance of  $(2X + 7) =$

- a. 25 b. 9 c. 36 d. 18 e. 43

$4 \text{Var } X = 36$

16. Standard deviation of  $X =$

- a. 4 b. 1 c. 7 d. 2 e. 3

$\sqrt{\text{Var } X} = \sqrt{9} = 3$

For 100 independent plays of the lottery define  $T = X_1 + \dots + X_{100}$ .

17.  $\text{Var } T =$

- a. 900 b. 300 c. 20 d. 140 e. 53

$\text{INDEP } 100 \text{Var } X = 900$

18.  $E T =$

- a. 30 b. 7 c. 200 d. 140 e. 3

$100 E X = 200$

19. Approximate 95% interval for  $T =$

- a. [120, 280] b. [180, 220] c. [170, 230] d. [135, 265] e. [140, 260]

$200 \pm 2\sqrt{900} = 200 \pm 60$

20-21. The distribution of IQ is normal with mean 100 and standard deviation 15.

20. The probability of IQ outside the range [100-30, 100+30] is approximately:

- a. 0.34 b. 0.05 c. 0.11 d. 0.68 e. 0.95 (draw a picture and work with pieces)

$1 - 0.9 = 0.05$

21. The probability of an IQ in range [100-15, 100] is approximately:

- a. 0.34 b. 0.05 c. 0.11 d. 0.68 e. 0.95



$\frac{1}{2}$  of 0.05